

## REVIEW

# Microbial ecology of organic aggregates in aquatic ecosystems

Meinhard Simon<sup>1,\*</sup>, Hans-Peter Grossart<sup>1</sup>, Bernd Schweitzer<sup>1</sup>, Helle Ploug<sup>2,\*\*</sup>

<sup>1</sup>Institute for Chemistry and Biology of the Marine Environment, University of Oldenburg, 26111 Oldenburg, Germany

<sup>2</sup>Marine Biological Laboratory, University of Copenhagen, Strandpromenaden 5, 3000 Helsingør, Denmark

**ABSTRACT:** Macroscopic organic aggregates, which are >500 µm and known as marine and lake snow, are important components in the turnover, decomposition and sinking flux of both organic and inorganic matter and elements in aquatic ecosystems. They are composed of various organic and inorganic materials depending largely on the given system and environmental conditions. The systems include the pelagic limnetic, the neritic and oceanic marine region, as well as shallow turbid environments, e.g. rivers, the littoral zone of lakes, estuaries and tidally affected coastal areas with intense turbulence and a high load of suspended matter. Aggregate abundance and size vary greatly among these systems. Macroaggregates are heavily colonized by bacteria and other heterotrophic microbes and greatly enriched in organic and inorganic nutrients as compared to the surrounding water. During the last 15 yr, many studies have been carried out to examine various aspects of the formation of aggregates, their microbial colonization and decomposition, nutrient recycling and their significance for the sinking flux. They have been identified as hot-spots of the microbial decomposition of organic matter. Further, microaggregates, which are <5 to 500 µm in size and stained by different dyes, such as transparent exopolymer particles (TEP) and Coomassie blue-stained particles, have been discovered and shown also to be important in the formation and decomposition of macroaggregates. In this review we give an overview of the present state of the microbial ecology of macro- and microaggregates, including the mentioned points but highlighting in particular the recent findings on the bacterial colonization of aggregates using molecular tools, their microbial decomposition and mineralization, and the significance of protozoans and metazoans for the colonization and decomposition of macroaggregates. Today it is evident that not only the aggregates but also their surroundings are sites and hot-spots of microbial processes, with the plume of solutes leaking out of the aggregates and greatly extending the volume of the intense decomposition processes. This microheterogeneity has important implications for the spatial and temporal dynamics of the organic-matter field in aquatic ecosystems and for our understanding of how heterotrophic organisms are involved in the decomposition of organic matter. The significance of aggregate-associated microbial processes as key processes and also for the overall decomposition and flux of organic matter varies greatly among the various systems, and is greatly affected by the total amount of suspended particulate matter. A conclusion from the presented studies and results is that the significance of bacteria for the formation and decomposition of aggregates appears to be much greater than previously estimated. For a better understanding of the functioning of aquatic ecosystems it is of great importance to include aggregate-associated processes in ecosystem modeling approaches.

**KEY WORDS:** Aggregates · Transparent exopolymer particles · Particulate organic matter · Algae · Bacteria · Zooplankton · Sinking flux · Dissolved organic matter

—Resale or republication not permitted without written consent of the publisher—

## 1. INTRODUCTION

One of the major topics in aquatic microbial ecology deals with the fate of particulate organic matter (POM) and the pathways by which it is (1) produced biologically or abiologically, (2) transferred within the food web or (3) transported downward as sinking POM through the water column, and decomposed and mineralized by microbes. POM interacts in various ways with colloidal and dissolved organic and inorganic materials, e.g. by adsorption or dissolution, and leads to the redistribution of elements, nutrients and specific compounds such as organic pollutants, heavy metals and radioisotopes in aquatic ecosystems. Operationally, the colloidal and dissolved material is separated from the particulate material by the fact that the latter is retained on filters with a pore size of 0.2 to 0.45  $\mu\text{m}$ . The primary and most important process of POM formation is photoautotrophic fixation of  $\text{CO}_2$  into biomass, in pelagic ecosystems usually by phytoplankton, but in coastal areas, shallow lakes, rivers and estuaries also by macrophytes, macroalgae and benthic microalgae. Hence, POM is initially living biomass of primary producers that, by various processes and after transfer steps in the food web, eventually turns into dead (detrital) POM often dominating the total POM. One important feature of organic particles is that they are densely colonized by various microbes, which often have key functions in the formation and decomposition of these detritus-dominated micropatches. The often very specific environmental conditions within these micropatches, distinctly different from the surrounding water, and the strikingly high activities of the particle-associated microbes, make them a unique and important subject in aquatic microbial ecology. Because many of these structures are composed of

smaller primary components, they are actually composites of detrital particles rich in organic matter and thus are called organic aggregates. Their macroscopic forms, i.e.  $>500 \mu\text{m}$ , are known as marine snow and lake snow aggregates (Fig. 1). These terms, as well as microaggregates, i.e. those  $<500 \mu\text{m}$ , include not only the detrital organic matter but also the associated microbes as integral components.

POM is one important component of the total suspended matter, consisting of living biomass, detrital organic and inorganic matter. The general significance of suspended matter in marine, estuarine, lacustrine and riverine ecosystems has been reviewed recently (Eisma 1993, Wotton 1994) but the microbial ecology of organic aggregates only to a lesser extent (Alldredge & Silver 1988, Alldredge 1992, Kirchman 1993, Lampitt 1996, Kjørboe 2001). These reviews, except the most recent one by Kjørboe (2001), do not include the recent progress in analyzing their decomposition and turnover of various substrates, the structural composition of aggregate-associated bacterial communities by molecular methods, and interactions with the surrounding water. However, the significance of POM, its microbial decomposition and its role in recycling of nutrients have been acknowledged and studied for more than 30 yr (Johannes 1968, Melchiorri-Santollini & Hopton 1972, Eppley et al. 1977, Cole et al. 1984, Kirchman & Ducklow 1987). The overall, but also very specific, significance of organic aggregates for the cycling and flux of elements and energy in pelagic marine, and more recently in lacustrine and estuarine ecosystems, in the light and significance of the microbial loop has emerged more and more only during the last 10 to 20 yr. Inspired by new concepts of the significance of aggregate-associated bacteria

for the cycling and decomposition of organic aggregates (Azam & Cho 1987, Cho & Azam 1988, Azam & Smith 1991, Smith et al. 1992, Grossart & Simon 1993) and by new technologies and methods to study the occurrence, size, spatio-temporal dynamics and microbial colonization of organic aggregates, research on this topic was boosted in the 1990s and led to the present knowledge in this field. In this review we focus on the formation and occurrence of organic aggregates and on heterotrophic processes of the associated microbes. Autotrophic processes are included only when appropriate.




Fig. 1. Size spectra of the major particulate and dissolved organic constituents in aquatic systems

## 2. METHODS TO STUDY AGGREGATE ABUNDANCE AND FORMATION

### 2.1. *In situ* abundance

Aggregate abundance *in situ* has been studied by SCUBA divers' visual observations, and by various types of still camera systems, also moored to sediment trap devices (e.g. Honjo et al. 1984, Johnson & Wangersky 1985, Asper 1987, Alldredge & Silver 1988, Costello et al. 1989, Alldredge 1991). Later, remotely or automatically controlled underwater still- and video-camera systems were applied to record vertical and temporal dynamics of aggregates in various size classes (e.g. Walsh & Gardner 1992, Lampitt et al. 1993a,b, Milligan 1995, Lampitt 1996, Jackson et al. 1997, Pilskaln et al. 1998). In combination with computer-assisted image analysis, these methods made it possible to establish *in situ* size spectra of aggregates and their dynamics, and resulted in new insights into their formation and short and long term dynamics.

### 2.2. *In situ* collection of macroaggregates

Macroscopic aggregates are very fragile and easily collapse when sampled by classical sampling devices such as Niskin and GoFlo bottles and when transferred into other bottles for further investigations. Therefore, special care has to be taken to sample these aggregates for further experiments. Still the best and most careful sampling method in pelagic environments is by SCUBA diving and the collection of macroscopic aggregates *in situ* in open-ended plastic syringes or widely opened bottles (Alldredge & Gotschalk 1990, Alldredge 1991). Once collected, aggregates remain in the syringes until further processing aboard the ship. Often it is unavoidable that aggregates in the syringes become more compact than when freely floating in the water. This collection method, however, limits aggregate studies to the upper 20 to 50 m. For sampling deeper layers, submarines and large volume transparent bottles were used (Alldredge & Youngbluth 1985, Lampitt et al. 1993a, Turley & Mackie 1994, Lampitt 1996). From these large bottles, individual aggregates can be collected visually once the bottle is hauled back on deck.

Sediment traps have been used for many years to collect sinking POM which is dominated by macroaggregates. Even though the elemental and biopolymer composition of this material can be examined (e.g. Cowie & Hedges 1992, Haake et al. 1992, Wakeham & Lee 1993), the aggregate structure is destroyed due to the long (days to years) deployment periods. Traps, exposed for short periods of time (hours), however,

were successfully used to collect fresh aggregates for studies of their bacterial colonization in which the aggregate size structure can be neglected (Grossart & Simon 1998a, Schweitzer et al. 2001).

Horizontal tubes and wide-mouthed funnels have been used to collect highly abundant aggregates in a very turbid estuarine environment, allowing the collection and gentle concentration of aggregates for further investigations (Kerner & Krogmann 1994, Zimmermann 1997). Possible biases by further aggregation and restructuring of the size distribution during the concentration procedure need careful examination.

### 2.3. Experimental formation of aggregates

Formation of macroscopic organic aggregates was first examined experimentally by Shanks & Edmondson (1989) in plexiglass cylinders rolling horizontally. This method gave the basis for many experimental studies examining details of the formation of aggregates and their microbial colonization and decomposition. This device is supposed to mimic the collision by differential settling of aggregates *in situ* and has often been used in various sizes and modifications since its introduction (Gotschalk & Alldredge 1989, Riebesell et al. 1991, Weiss et al. 1996, Artolozaga et al. 1997, Grossart et al. 1997, 1998, Unanue et al. 1998a, Knoll et al. 2001). Because the hydrodynamics in the rolling cylinders are somewhat different from *in situ* conditions with enhanced shear during the start-up phase, Couette flocculators were suggested as a better means to mimic *in situ* hydrodynamics (Jackson 1994) and were used in several studies (e.g. Kiørboe & Hansen 1993, Drapeau et al. 1994). Aggregates are more difficult to collect in Couette flocculators, which limits the use of these devices. Vertically rotating glass syringes containing only 1 or a few macroaggregates were used to examine autotrophic and heterotrophic activities and colonization, and interactions with the surrounding water (Biddanda 1988, Biddanda & Pomeroy 1988, Gotschalk & Alldredge 1989, Grossart & Simon 1998b, Schweitzer et al. 2001).

## 3. OCCURRENCE OF ORGANIC AGGREGATES

### 3.1. Size, abundance and fractal dimension

In pelagic systems, phytoplankton primary production forms biological particles in the size range of 1  $\mu\text{m}$  (picoplankton cells) to a few hundred micrometers, and in extreme cases a few millimeters (diatom chains) or centimeters (cyanobacterial filaments), depending

on the trophic state, seasonal phase and geographic region of the system. Yet, organic particles and aggregates occur in the size range of  $<1\ \mu\text{m}$  to beyond 10 cm and thus cover a much wider range than biological particles—more than 6 orders of magnitude (Fig. 1), implying that many complex physical, chemical, biological and specific microbial processes are involved in the formation and decomposition of organic particles and aggregates. In systems dominated by macroalgae, macrophytes or allochthonous inputs, even larger particles are the primary sources of organic aggregates. Various processes contribute to the break-down of these large particles, such as mechanical disruption, consumption by detritivores, and microbial decomposition. There is a continuum in the size distribution but special types of aggregates often occur in distinct size classes, which often reflect their origin, significance and fate. Operationally, 3 size classes of organic aggregates are distinguished (Fig. 1): macroscopic aggregates  $>500\ \mu\text{m}$  (macroaggregates, marine snow, lake snow), microscopic aggregates 1 to  $500\ \mu\text{m}$  (microaggregates), and submicron particles  $<1\ \mu\text{m}$ , discovered a decade ago (Koike et al. 1990, Wells & Goldberg 1991). There is no evidence that submicron particles are involved in the formation or break-down of micro- and macroaggregates. The abundances of the smallest microparticles and largest aggregates range from  $<1$  to  $>10^8\ \text{l}^{-1}$  and are inversely related to size.

It has been shown that the structure of macro- and microaggregates is best described by fractal geometry (Logan & Wilkinson 1990, Kilps et al. 1994, Chen & Eisma 1995). Exceptions are abandoned larvacean houses, which are not formed by aggregation (Alldredge 1998). Fractal geometry means that the mass or number of component particles ( $N$ ) and the characteristic length scale of the aggregate ( $l$ ) are related according to the function  $N \sim l^{D_n}$ .  $D_n$  ( $D_1$ ,  $D_2$ ,  $D_3$ ), a non-integer between 1 and 3, is the fractal dimension of the aggregate and a function of the perimeter ( $P \sim D_1$ ), the area ( $A \sim D_2$ ) or the volume ( $V \sim D_3$ ). Only when  $D_n$  is an integer does the aggregate have an Euclidian geometry, e.g.  $D_3 = 3$  for a sphere. Marine snow typically has a fractal dimension of  $D_3$  between 1.3 and 2.1, which is much lower than that of inorganic colloidal aggregates and only slightly higher than that of activated sludge flocs (Logan & Wilkinson 1990, Logan & Kilps 1995, Jackson et al. 1997).  $D_1$  and  $D_2$  of aggregates in the shallow North Sea and the Elbe estuary, Germany, range between 1.01 and 1.14, and between 1.41 and 1.81, respectively (Chen & Eisma 1995), and  $D_2$  of marine snow of various types in Californian coastal waters between 1.1 and 1.8 (Kilps et al. 1994). Fractal geometry allows estimation of the porosity of aggregates, i.e. pore volume, which is inversely related to the fractal dimension (Logan & Wilkinson

1990). The porosity is important in controlling the aggregate's sinking rate, the flux of water through the aggregate moving relative to the surrounding water, and the flux of nutrients and substrates to and from the microorganisms colonizing the aggregate's surface (Logan & Hunt 1987, Alldredge & Gotschalk 1988, Logan & Alldredge 1989, Ploug 2001).

### 3.2. Formation of organic aggregates

Aggregates are formed by a variety of physical, physico-chemical and biological processes. In particular, the former have been studied and reviewed quite extensively (e.g. O'Melia 1987, Eisma 1993, Johnson et al. 1994, Stumm et al. 1994). These reviews provide the basis for the outlines below. Biological factors and the significance of microorganisms in aggregation processes have attracted detailed interest only fairly recently. Generally, aggregation is a complex process controlled by 3 primary variables: (1) the particle concentration, density, size distribution and shape; (2) shear and differential settling velocities; and (3) the probability with which the particles stick together after collision. The sticking coefficient ( $\alpha$ ) describes the ratio of numbers of successful collisions/numbers of total collisions (McCave 1984). It varies considerably for different types of marine macroaggregates and ranges from  $<0.2$  for non-sticky material to 0.60–0.88 for sticky phytodetrital aggregates (Alldredge & McGilvary 1991, Kjørboe et al. 1994, 1996, Waite et al. 1997). Interestingly, the stickiness of diatoms and aggregates appears to be highest during the transition from the exponential growth phase to the stationary phase and decreases again thereafter (Kjørboe et al. 1990, Kjørboe & Hansen 1993, Drapeau et al. 1994, Dam & Drapeau 1995). In contrast, Engel (2000) reported an increasing stickiness during the decline of an experimental diatom bloom and highest values at the end. Transparent exopolymer particles (TEP) have been identified as very important in controlling the sticking coefficient of diatoms and other organic particles (Alldredge et al. 1993, see Section 3.4.1). The following mechanisms causing collisions of suspended particles have been identified:

**Brownian motion:** Collision of particles due to random walk. It is only important for particles  $<8\ \mu\text{m}$  because their movement is largely controlled by laminar flow and their diffusivity (Eisma 1993).

**Shear:** Particles in laminar or turbulent shear collide if the distances of their flow streamlines or eddies are smaller than the sum of the particle radii. It is important for the collision of particles  $>8\ \mu\text{m}$  (Eisma 1993) and is the dominant mechanism at interfaces such as at discontinuity layers in the water column, in the bottom

nepheloid layer, or at tidal currents in estuaries and shallow seas. In pelagic systems, shear is one of the major factors controlling aggregation (Jackson 1990). At energy dissipation rates of  $10^{-7}$  to  $>10^{-4}$   $\text{m}^2 \text{s}^{-3}$ , which can occur at the abovementioned discontinuity layers, the resulting shear may lead to disaggregation or size reduction of macroaggregates, depending on the physical strength and size of the aggregates (Alldredge et al. 1990, Riebesell 1991a, 1992, McIntyre et al. 1995).

*Differential settling:* Settling particles can overtake, and intercept, more slowly sinking particles. As with shear, particles collide if the distances of their flow streamlines are smaller than the sum of the particle radii. Usually, larger, more rapidly sinking particles scavenge smaller particles by this mechanism (Jackson 1990, Kepkay 1994). If the larger particle is surrounded by a diffusive boundary layer, the collision also depends on the diffusion rate of the smaller particle. This may be important for scavenging submicron particles and colloids. Differential settling is important for sinking particles in the water column and in slack water in tidally affected shallow seas and estuaries.

*Filtration:* Theoretical studies of fluid dynamics have shown that highly porous aggregates may experience an interstitial water flow in the order of tenths to hundreds of  $\mu\text{m s}^{-1}$  when the aggregate is sinking (Logan & Hunt 1987, Logan & Alldredge 1989) such that nutrients, but also colloids or other particles, are collected by size exclusion (Stolzenbach 1993). Interstitial fluid flow through aggregates, however, has not been experimentally quantified due to insufficient spatial resolution and sensitivity of existing techniques. The high content of mucus and TEP within aggregates may in fact limit advection through aggregates. It is not known yet how important this mechanism is in aggregate formation. It may be important under certain conditions for passively capturing particles that do not actively form aggregates such as discarded larvacean houses, pteropod webs and non-sticky dinoflagellates (Alldredge & Silver 1988, Alldredge et al. 1998, Silver et al. 1998).

*Biological capture:* Zooplankton capture other particles by filtering food particles and thus aggregate such particles in the filtering apparatus or the gut. This process may contribute to structuring the composition of aggregates consisting of abandoned larvacean houses, pteropod webs and fecal pellets (Pomeroy & Deibel 1980, Lampitt et al. 1990, Bochdansky & Herndl 1992, Herndl 1992).

Whether the collisions are successful or not depends on the following factors:

*Surface properties of the particle:* A particle or aggregate has attractive or repulsive surface properties, depending on its size, composition and chemical

coating. Most particles are negatively charged at the surface because of the adsorbed organic material with many carboxylic or hydroxylic compounds (Gibbs 1983, O'Melia 1987, Johnson et al. 1994, Stumm et al. 1994). In the presence of coatings of organic multilayers, particles may also exhibit hydrophobic attractive properties (Zutic & Tomaic 1988). Different surface properties of individual diatom species, depending also on their physiological state and the microbial colonization, lead to selective aggregation and removal of diatoms during phytoplankton blooms (Riebesell 1991b, Passow et al. 1994, Alldredge et al. 1995, 1998, Hansen et al. 1995, Hansen & Kjørboe 1997).

*Bridging with divalent cations:* Divalent cations such as  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  act as chemical bridging agents enhancing the attraction and sticking properties of electronegatively charged particles (Eisma 1993, Stumm et al. 1994). They are most important in mediating aggregation of electronegatively charged particles and are the main reason why aggregation is a positive function of salinity.

*Mediation of coagulation and aggregation by dissolved organic matter:* Dissolved organic matter (DOM) may enhance or reduce aggregation depending on its quality and concentration (Johnson et al. 1994). DOM, and in particular humic substances with a high negative charge, readily adsorb to particles but also reduce aggregation in the absence of cations such as  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  (e.g. Gerritsen & Bradley 1987, Zutic & Tomaic 1988, Eisma 1993). Therefore, aggregation can be expected to be negatively correlated with DOM concentration, if DOM is dominated by dissolved humics, and to be lowest in humic lakes. On the other hand, low concentrations of DOM enhance aggregation of inorganic particles exhibiting a positive surface charge when not coated by organic material (Gibbs 1983, Johnson et al. 1994). DOM of high molecular weight is more effective in surface-reactive properties than DOM of low molecular weight. Dissolved or particulate organic polymers produced by phytoplankton and bacteria have also been shown to be important in enhancing the formation of macroaggregates and in stabilizing the structure of aggregates (see Section 3.4.1).

There are also other processes resulting in aggregation:

*Bubbling (surface coagulation) of DOM:* Surface-reactive DOM, mainly polysaccharides, can be converted into particles through coagulation at the surfaces of gas bubbles when they burst or collapse (e.g. Johnson et al. 1986, Eisma 1993, Mopper et al. 1995, Zhou et al. 1998, Mari 1999). Even though this process has been studied quite extensively and occurs during the development of diatom blooms, it is not yet clear to what extent it contributes to the formation of aggregates. Presumably it is important under certain condi-



tions, such as at the sea surface, in the formation of TEP (see Section 3.4.1).

*Spontaneous coagulation of DOM:* DOM can form colloidal material, submicron particles and eventually also microaggregates due to spontaneous coagulation in the dissolved phase (Kepkay 1994, Chin et al. 1998, Mari & Burd 1998, Passow 2000). The occurrence of this process has been demonstrated convincingly but it is not yet known to what extent it contributes to the formation of aggregates. Most recent results suggest that it is important under certain conditions in the formation of TEP (Passow 2000, Section 3.4.1).

Biological processes may also directly result in and control the formation of aggregates by various mechanisms:

*Larvaceans discarding their houses:* Larvaceans discarding their houses with a clogged filtration apparatus directly generate macroaggregates because living and senescent algae, protozoans, bacteria and microaggregates are attached to these houses. In fact, abandoned larvacean houses are one of the most abundant types of marine snow in certain areas (Alldredge & Madin 1982, Alldredge & Silver 1988, Hansen et al. 1996, Silver et al. 1998).

*Detached pteropod nets:* Pteropods can lose their filter nets when they become clogged by various particles such as algal cells and microaggregates and thus directly produce macroaggregates in pelagic marine environments.

*Defecation and molting of zooplankton:* All marine and most lacustrine zooplankton produce fecal pellets, which consist of densely packed micro- or macroaggregates surrounded by a peritrophic membrane. These fecal pellets can stick to other aggregates or directly sink as individual particles to deeper layers in the water column, thus contributing substantially to the formation of sinking aggregates and sinking flux (e.g. Lampitt et al. 1990, Bochdansky & Herndl 1992, Turner 2002). Only herbivorous cladocerans in lakes, such as daphnids, do not produce fecal pellets and thus are differently involved in the formation of macroaggregates. This notion has implications for differences in the relationship between primary production and the sinking flux in marine and cladoceran-dominated lacustrine ecosystems (Aksnes & Wassmann 1993). However, due to the short generation time of their parthenogenetic generations (4 to 10 d), cladocerans can establish very high abundances for short periods of time and produce high amounts of molts and carcasses, which may dominate sinking macroaggregates in lakes for short periods of time (Grossart et al. 1997, Gries & Güde 1999). Holo- and meroplanktonic zooplankton may also contribute substantially to sinking macroaggregates in shallow bays (Shanks & Edmondson 1990, Shanks & Del Carmen 1997, Shanks & Walters 1997, see Section 4.1).

*Mediation of aggregation by microbes:* There is growing evidence demonstrating the direct significance of microorganisms in the formation of aggregates and that aggregation is a direct consequence of the activity of microorganisms: (1) Biddanda (1985) experimentally showed that the formation of marine macroaggregates was only possible in the presence of live microorganisms because no aggregation occurred when the samples were poisoned by mercury chloride or when bacteria were removed by filtration. (2) Paerl (1974) demonstrated by autoradiography that aggregate-associated bacteria take up DOM of low molecular weight from the surrounding water and convert it to fibrillar material gluing together microaggregates. Other authors also demonstrated the secretion of fibrillar material by bacteria, leading to strengthened aggregate structure (Leppard et al. 1979, Heissenberger et al. 1996). (3) Decho (1990), Alldredge et al. (1993) and thereafter many other authors showed that diatoms secrete large amounts of acidic mucopolysaccharides, named TEP, which are very important in and often control the formation of macroaggregates in pelagic ecosystems (see Section 3.4.1). Also, bacteria were recently shown to secrete large amounts of TEP and thus may directly control the formation of macroaggregates (Stoderegger & Herndl 1999). (4) Specific bacterial populations colonizing diatoms or living in their phycosphere (Bell & Mitchell 1972) obviously also control phytoplankton growth, mortality and secretion of polysaccharides, and thus may directly control the aggregate formation (Azam 1998, Brussaard & Riegman 1998, Guerrini et al. 1998, Azam et al. 1999, Grossart 1999, Herndl et al. 1999).

The net result of aggregation is a function of all the abovementioned parameters and processes, varying largely according to the type of aggregate, the environment and the organisms involved. There is growing evidence that phytoplankton and bacteria are often directly involved in and even may control aggregate formation. But also when aggregation may be a non-biological process, microorganisms indirectly affect it by secreting and consuming organic material, which leads to or prevents aggregation.

### 3.3. Macroscopic aggregates

Macroscopic aggregates were discovered in the 1950s by submarines and called marine snow due to their snowflake-like appearance when reflecting an underwater light beam (Fig. 2A; Suzuki & Kato 1953). They have been studied extensively in marine systems since then and more recently, to a lesser extent, also in lakes, rivers and estuaries (Fowler & Knauer 1986, Alldredge & Silver 1988, Alldredge & Gotschalk 1990,




Fig. 2. *In situ* photograph of a marine snow aggregate (A) from a pelagic environment and (B) of microaggregates in a shallow turbid environment (photo M. Lunau)

Allredge 1992, 1998, Herndl 1992, Grossart & Simon 1993, 1998a,b, Berger et al. 1996, Lampitt 1996, Grossart et al. 1997, 1998, Zimmermann 1997, Ploug et al. 1999, Grossart & Ploug 2000, Neu 2000). The term 'lake snow' was adopted from marine snow for limnetic macroaggregates (Grossart & Simon 1993) and most recently 'river snow' for riverine aggregates (Neu 2000). Coagulation theory has also been applied to aggregate formation (e.g. Jackson 1990, 1995, Hill 1992, Hansen et al. 1995). Today, the general ecological significance of macroaggregates is fairly well understood. In most oceanic and neritic environments and in the lakes studied so far, their size ranges between 300 and 500  $\mu\text{m}$  to several centimeters, and their abundance between  $<1$  and  $\sim 100 \text{ l}^{-1}$  (Table 1). Macroaggregates in the pelagic zone of lakes appear to be smaller than those in pelagic marine environments. The largest aggregates of  $\sim 20 \text{ cm}$  and of 1 to 4 m in size were reported from the mesopelagic zone off California (Steinberg et al. 1997) and from the Northern Adriatic Sea (Herndl 1992, Degobbis et al. 1999, Herndl et al. 1999), respectively. The large aggregates in the Northern Adriatic Sea exhibited a very dense structure, and their appearance was described as gel-like or as a false benthos (Kaltenböck & Herndl 1992). Diatom aggregates of  $\sim 20 \text{ cm}$  in size were also observed in Danish coastal waters (H.-P. Grossart unpubl. results) and may be more abundant than assumed. Due to the fragile nature of these large aggregates they are very difficult to sample.

Across systems, the abundance of macroaggregates is positively related to phytoplankton biomass (Allredge & Gotschalk 1990), but on a temporal and vertical scale, abundances of macroaggregates often lag behind phytoplankton growth and peak at the end of

blooms, with highest numbers below the chlorophyll maximum (Allredge & Gotschalk 1989, Riebesell 1991b, 1992, Lampitt et al. 1993a, Kjørboe et al. 1994, Allredge et al. 1995, Grossart et al. 1997, Walsh et al. 1997, Becquevort & Smith 2001). In fact, diatom blooms often, but not always, are terminated by aggregation events and sink out of the mixed layer (Smetacek 1985, Kjørboe et al. 1994, 1996, 1998, Allredge et al. 1995, Logan et al. 1995, Tiselius & Kuylenstierna 1996). Thus, macroaggregates dominate the sinking POM in marine as well as in lacustrine systems (Fowler & Knauer 1986, Asper 1987, Allredge & Silver 1988, Riebesell 1991a, Walsh et al. 1997, Grossart & Simon 1998a, Pilskaln et al. 1998). Sinking rates of macroaggregates range from  $<5$  to  $200 \text{ m d}^{-1}$  and are a positive function of their size (Allredge et al. 1987, Asper 1987, Allredge & Gotschalk 1988, 1989, Allredge & Silver 1988, Grossart & Simon 1993, Kjørboe et al. 1994, Hansen et al. 1996, Pilskaln et al. 1998). Sinking rates determined *in situ*, however, were lower than those determined in settling chambers (Allredge et al. 1987, Allredge & Gotschalk 1988), presumably because aggregates become more compact when collected in syringes and transferred to the containers in which sinking rates are determined experimentally. Further, net sinking rates *in situ* are also the results of turbulent water movements and orbital motions, which are most pronounced close to the surface. In fact, aggregates can persist in the mixed layer longer than assumed from their sinking rates (McIntyre et al. 1995) and as reflected by the degradation state of the associated fecal pellets (Allredge et al. 1987). At density discontinuity layers, sinking aggregates may accumulate because of a reduced excess density or enhanced shear rates (Allredge & Crocker 1995, McIntyre et al.

Table 1. Abundances of macroaggregates and microaggregates in various aquatic environments. Macroaggregates were quantified visually, or by underwater (UW) photo- or videography. CSP: Coomassie-stained particles; DAPI: 4',6-diamidino-2-phenylindole-stained particles; TEP: transparent exopolymer particles

Location	Depth (m)	Macroaggregates l <sup>-1</sup>	Microaggregates l <sup>-1</sup>	Type	Source
Southern California Bight	7–15	<1–8			Allredge (1979)
Gulf of California	7–15	<1–3.7			Allredge (1979)
Northern Atlantic	7–15	0.1–1.0			Allredge et al. (1986)
Southern California Bight	7–15	1.4–4.0			Allredge et al. (1986)
Subtropical NW Atlantic	120–530	<0.04			Allredge & Youngbluth (1985)
Southern California Bight	7–15	0.2–1.7			Allredge & Gotschalk (1990)
Monterey Bay	100–500	0.5–50 <sup>a</sup>			Pilskaln et al. (1998)
Southern California Bight	7–15	<20–79.4			Graham et al. (2000)
Equatorial Pacific	0–80	10–55 <sup>a</sup>			Walsh et al. (1997)
Equatorial Pacific	100–1000	2–15 <sup>a</sup>			Walsh et al. (1997)
Northern Atlantic	270	<0.5–12 <sup>a</sup>			Lampitt et al. (1993b)
Benguela upwelling	0–30	85–640 <sup>a</sup>	5.5 × 10 <sup>6</sup> –16 × 10 <sup>6</sup>	TEP	Kjørboe et al. (1998)
Coastal Atlantic	0–15	13.5–62.8			Shanks & Del Carmen (1997)
Coastal Pacific	0–15	1–11			Shanks & Del Carmen (1997)
Kattegat (North Sea)	0–50	<130			Tiselius & Kuylenstierna (1996)
North Sea, German Bight	0–30	425–5300 <sup>a</sup>			Riebesell (1991a)
Elbe estuary	1	20–4000			Zimmermann (1997)
Baltic Sea (benthic boundary layer)	16–26	152–787			Jähmlich et al. (1999)
Lake Constance	5–25	<1–50			Grossart & Simon (1993)
Lake Constance	5–25	<1–50	0.5 × 10 <sup>2</sup> –1.5 × 10 <sup>6</sup>	TEP	Grossart et al. (1997)
Lake Constance	6–100		3.1 × 10 <sup>5</sup> –1.5 × 10 <sup>6</sup>	DAPI	Brachvogel et al. (2001)
Lake Constance	6–100		7.1 × 10 <sup>4</sup> –2.6 × 10 <sup>6</sup>	TEP	Brachvogel et al. (2001)
Lake Kinneret	15–35		1.0 × 10 <sup>5</sup> –7.0 × 10 <sup>6</sup>	TEP	Grossart et al. (1998)
Lake Kinneret	1–38		4.4 × 10 <sup>5</sup> –2.5 × 10 <sup>7</sup>	TEP	Berman & Viner-Mozzini (2001)
Lake Kinneret	1–38		1.4 × 10 <sup>5</sup> –1.2 × 10 <sup>7</sup>	CSP	Berman & Viner-Mozzini (2001)
Frederiksborg Slotssø	0.5		0.84 × 10 <sup>8</sup> –8.7 × 10 <sup>8</sup>	TEP	Worm & Søndergaard (1998b)
Southern California Bight	10		2.8 × 10 <sup>4</sup> –4.0 × 10 <sup>5</sup>	TEP	Allredge et al. (1993)
Monterey Bay	5–100		1.5 × 10 <sup>5</sup> –4.9 × 10 <sup>6</sup>	TEP	Allredge et al. (1993)
Monterey Bay	5–76		2.5 × 10 <sup>5</sup> –5.2 × 10 <sup>6</sup>	TEP	Passow & Allredge (1994)
Southern California Bight	10		2.5 × 10 <sup>4</sup> –6.3 × 10 <sup>5</sup>	TEP	Passow & Allredge (1994)
Subtropical NW Atlantic	300–1400		2.0 × 10 <sup>3</sup> –6.0 × 10 <sup>3</sup>	TEP	Passow & Allredge (1994)
Kattegat (North Sea/Baltic Sea)	0–25		3.0 × 10 <sup>6</sup> –6.0 × 10 <sup>7</sup>	TEP	Mari & Kjørboe (1996)
Kattegat (North Sea/Baltic Sea)	0–30		5.0 × 10 <sup>7</sup> –3.8 × 10 <sup>8</sup>	TEP	Mari & Burd (1998)
Northern Adriatic Sea	0–15		<1.0 × 10 <sup>3</sup> –6.0 × 10 <sup>5</sup>	TEP	Schuster & Herndl (1995)
NW Mediterranean Sea	0–40		1.0 × 10 <sup>7</sup> –2.2 × 10 <sup>8</sup>	TEP	Mari et al. (2001)
NW Mediterranean Sea	0–75		2.2 ± 3.2 × 10 <sup>6</sup>	DAPI	Mostajir et al. (1995)
Scripps Pier, Southern California Bight	5		2.0 × 10 <sup>7</sup> –3.4 × 10 <sup>7</sup>	CSP	Long & Azam (1996)
Arabian Sea	0–1500		2.0 × 10 <sup>6</sup> –3.5 × 10 <sup>7</sup>	CSP	Long & Azam (1996)

<sup>a</sup>UW photography

1995). In fact, there is good evidence that the initial accumulation of mucilage at the density discontinuity layer in the northern Adriatic Sea is a result of this phenomenon (Allredge 1999, Degobbi et al. 1999, Herndl et al. 1999). Aggregates dominated by cyanobacteria-derived material exhibit very low sinking rates or do not sink at all due to the very low excess density (Grossart et al. 1997). Macroaggregates with intensive primary production and entrapped oxygen gas bubbles may even move upward in the water column (Riebesell 1992, Herndl 1992). This may also occur when filamentous and gas vacuole-containing cyanobacteria are abundant (Capone et al. 1997, Stal et al. 1999).

In pelagic systems, numbers of macroaggregates generally decrease with depth, but the size does not. Due to the differential settling rates of aggregates of different sizes, larger aggregates accumulate with depth (Allredge & Gotschalk 1989). Diurnal differences in the aggregate abundance with highest numbers during the day were reported from the mixed layer (Graham et al. 2000) as well as the upper mesopelagic zone (Lampitt et al. 1993b, Walsh et al. 1997), presumably due to the diurnal vertical migration of zooplankton affecting aggregation in various ways (see below). Possible effects of light-dependent formation of TEP and mucus by phytoplankton or enhanced microbial decomposition during the night need be examined in future studies.



In estuaries and tidally affected shallow seas, macroaggregates are usually smaller than in lakes and the open sea with a size of  $<2000\ \mu\text{m}$  due to high shear rates leading to disaggregation (Chen et al. 1994, Chen & Eisma 1995, Riebesell 1991a, 1992, Zimmermann 1997, Zimmermann-Timm et al. 1998). In the Elbe estuary, pronounced seasonal variations with smaller aggregates in fall and winter and a continuously decreasing size toward the sea occur (Zimmermann 1997, Zimmermann-Timm et al. 1998). Abundances in these systems are much higher than in lakes and the open sea due to high resuspension rates (Table 1), and highly dynamic changes and redistributions in abundance and size occur within tidal cycles (Eisma 1993, Eisma & Li 1993, Milligan 1995, Zimmermann 1997). This is particularly true for the turbidity maximum, which is established in tidally affected estuaries at salinities between 2 and 6‰ and has great implications for physico-chemical and microbial transformation of dissolved and particulate matter (Eisma 1993, Crump & Baross 1996, Förstner 1996, Grabemann et al. 1996, Crump et al. 1998).

Also in the benthic boundary layer, macroaggregates have been observed even though very little information about their formation and composition is available. Preliminary results from the Baltic Sea show that their abundance is higher than in the pelagic zone (Table 1) and their size smaller (Jähmlich et al. 1999). Because high shear, intense resuspension and thus differential settling of particles occur, aggregate formation presumably is an important phenomenon in this environment, which needs to be studied more intensively. According to reports on the fluff layer in the deep ocean, macroaggregates also play an important role in structuring the benthic boundary layer in this environment (Lochte & Turley 1988, Thiel et al. 1988, Thomsen & Graf 1994).

### 3.3.1. Composition

Macroaggregates are composed of component particles consisting of living, senescent and dead algae, mainly diatoms, but also of coccolithophorids, cysts of thecate dinoflagellates, filamentous cyanobacteria, phytodetritus, diatom frustules, bacteria, protozoans, zooplankton molts and carcasses, abandoned larvacean houses, pteropod webs, fecal pellets, macrophyte detritus, clay and silt minerals, calcite and other particles scavenged from the surrounding water (Alldredge & Silver 1988, Alldredge & Gotschalk 1990, Riebesell 1991b, Herndl 1992, Kaltenböck & Herndl 1992, Grossart & Simon 1993, Lampitt et al. 1993a, Zimmermann & Kausch 1996, Grossart et al. 1997, Alldredge et al. 1998, Kiørboe et al. 1998, Zimmermann-

Timm et al. 1998). Most typical in pelagic marine systems are macroaggregates composed of abandoned larvacean houses, fecal material, diatoms and miscellaneous unidentifiable materials in aging systems. Cysts of dinoflagellates and empty thecae obviously can be scavenged by diatom aggregates and make the aggregates extra dense and rich in organic matter but do not actively form macroaggregates (Alldredge et al. 1998). Algae often are highly enriched on aggregates as compared to the surrounding water (Table 2). Cyanobacterial colonies can be important components of macroaggregates in tropical marine systems (Capone et al. 1997), the Baltic Sea (Stal et al. 1999) and stratified lakes (Grossart et al. 1997, 1998, Worm & Søndergaard 1998a). In shallow seas and estuaries and in the littoral zone of lakes, macrophyte debris, resuspended minerals and organic particles are important in the formation and redistribution of macroaggregates (Eisma 1993, Eisma & Li 1993, Chen & Eisma 1995, Milligan 1995, Grossart et al. 1998). Macroaggregates harbor a rich community of heterotrophic microbes, predominantly bacteria, which contribute only a minor fraction to the organic carbon bound in macroaggregates but are most important for their decomposition and undergo a pronounced succession (see below). Usually, the composition of the macroaggregates reflects the particle spectrum in the water column, even though quantitative differences may also occur due to selective aggregation properties of the primary particles (see above).

The dry weight of pelagic marine and limnetic macroaggregates ranges from 3 to  $>200\ \mu\text{g}$  per aggregate and is a positive function of the aggregate size (Alldredge & Gotschalk 1988, Alldredge 1998, Alldredge et al. 1998, Ploug & Grossart 2000). Organic carbon constitutes 10 to 40% of the total aggregate dry weight in marine and estuarine aggregates (Alldredge 1979, 1998, Eisma 1993) and up to 66% in limnetic aggregates, with highest percentages when cyanobacterial components are dominant (Grossart & Simon 1993, 1998b). Dissolved organic carbon (DOC) in the interstitial water of marine snow aggregates comprises 20% of the total organic carbon (particulate organic carbon [POC] + DOC) and is a non-negligible fraction of the organic carbon (Alldredge 2000). Dry weight, POC, DOC and particulate organic nitrogen (PON) significantly increase with the aggregate volume to the power of 0.49 to 0.54 (Alldredge 1998). The dry mass, however, occupies only  $<1\%$  of the total aggregate volume and its relative proportion decreases with the aggregate size. This relationship reflects the fractal nature of macroscopic aggregates and their high porosity, which increases with the aggregate size (Alldredge & Gotschalk 1988, Logan & Alldredge 1989, Logan & Wilkinson 1990, Alldredge 1998). The C:N (w:w) ratio ranges between 5:1 and 20:1 and is a posi-

Table 2. Colonization of macroaggregates (agg) by various organisms and enrichment factors (EF) relative to the surrounding waters. EF were calculated as the abundances of organisms per volume aggregate divided by the abundances per volume surrounding water. na: not available

Organism	Location	Cells (ind.) agg <sup>-1</sup>	EF	Source
<b>Algae</b>				
Phytoplankton <2 µm	NW Atlantic	na	19–43	Caron et al. (1986)
Phytoplankton 2–20 µm	NW Atlantic	na	7–67	Caron et al. (1986)
Phytoplankton >20 µm	NW Atlantic	na	13–700	Caron et al. (1986)
Diatoms	Monterey Bay	$5.2 \times 10^3$	<27	Silver et al. (1978)
Diatoms	Monterey Bay	$6.0 \times 10^4$ – $4.2 \times 10^5$	$130$ – $1.7 \times 10^3$	Silver et al. (1998)
Synechococcus	Monterey Bay	$1.9 \times 10^6$ – $5.4 \times 10^6$	45–160	Silver et al. (1998)
Diatoms	Southern California Bight	$4.1 \times 10^2$ – $6.0 \times 10^3$	$160$ – $3.9 \times 10^3$	Allredge et al. (1998)
Dinoflagellates	Southern California Bight	$0.53 \times 10^2$ – $1.2 \times 10^3$	$110$ – $1.2 \times 10^4$	Allredge et al. (1998)
Synechococcus	NE Atlantic	$1.0 \times 10^4$ – $2.5 \times 10^4$	$1.0 \times 10^3$ – $7.6 \times 10^3$	Turley & Mackie (1994)
<b>Zooplankton</b>				
Harpacticoid copepods	Northern Adriatic Sea	0–250	3.5–10	Bochdansky & Herndl (1992)
Nauplii	Northern Adriatic Sea	0–1233	3.7–145	Bochdansky & Herndl (1992)
Polychaete larvae	Northern Adriatic Sea	30–3633	344–765	Bochdansky & Herndl (1992)
Juvenile turbellarians	Northern Adriatic Sea	0–1208	62–670	Bochdansky & Herndl (1992)
Calanoid copepods	Gulf of Mexico	0.1–32.9	0.9–1.9	Green & Dagg (1997)
Cyclopoid copepods	Gulf of Mexico	0.6–42	0.8–100	Green & Dagg (1997)
Harpacticoid copepods	Gulf of Mexico	0.2–14.3	18.5–57.1	Green & Dagg (1997)
Crustacean nauplii	Gulf of Mexico	1.6–72.7	100	Green & Dagg (1997)
Polychaete larvae	Gulf of Mexico	0.1–11.4	7.1–35.8	Green & Dagg (1997)
Molluscs	Gulf of Mexico	0.2–1.4	0.2–12.2	Green & Dagg (1997)
Salps	Gulf of Mexico	0.2–1.4	0.4–6.9	Green & Dagg (1997)
Larvaceans	Gulf of Mexico	0.1–26.2	8.2–35.1	Green & Dagg (1997)
Ascidians	Gulf of Mexico	1.0–4.1	0.45	Green & Dagg (1997)
Calanoid copepods	Coastal Atlantic	na	<3 <sup>b</sup>	Shanks & Edmondson (1990)
Cyclopoid copepods	Coastal Atlantic	na	<3 <sup>b</sup>	Shanks & Edmondson (1990)
Harpacticoid copepods	Coastal Atlantic	na	2–35 <sup>b</sup>	Shanks & Edmondson (1990)
Crustacean nauplii	Coastal Atlantic	na	0.9–2 <sup>b</sup>	Shanks & Edmondson (1990)
Nematodes	Coastal Atlantic	na	50–100 <sup>b</sup>	Shanks & Edmondson (1990)
Polychaete larvae	Coastal Atlantic	na	3–20 <sup>b</sup>	Shanks & Edmondson (1990)
Polychaetes	Coastal Atlantic	na	60–87 <sup>b</sup>	Shanks & Edmondson (1990)
Polychaete larvae	Coastal Atlantic	<1	18–70 <sup>b</sup>	Shanks & Del Carmen (1997)
Polychaete larvae	Coastal Pacific	<1	3–27 <sup>b</sup>	Shanks & Del Carmen (1997)
Larvaceans	Coastal Atlantic/Pacific	na	2–3 <sup>b</sup>	Shanks & Walters (1997)
Calanoid/cyclopoid copep.	Coastal Atlantic/Pacific	na	0.1–2 <sup>b</sup>	Shanks & Walters (1997)
Harpacticoid copepods	Coastal Atlantic/Pacific	na	2–55 <sup>b</sup>	Shanks & Walters (1997)
Copepod nauplii	Coastal Atlantic/Pacific	na	1–11 <sup>b</sup>	Shanks & Walters (1997)
Nematodes	Coastal Atlantic/Pacific	na	8–88 <sup>b</sup>	Shanks & Walters (1997)
Foraminifera	Coastal Atlantic	na	47 <sup>b</sup>	Shanks & Walters (1997)
Bivalve veliger	Coastal Atlantic/Pacific	na	4.5–10 <sup>b</sup>	Shanks & Walters (1997)
Gastropod veliger	Coastal Atlantic/Pacific	na	2–34 <sup>b</sup>	Shanks & Walters (1997)
Anthozoan larvae	Coastal Pacific	na	2–32 <sup>b</sup>	Shanks & Walters (1997)
Total copepods	Coastal Pacific	109.8 ± 28.4	197	Steinberg et al. (1994)
Calanoid copepods	Coastal Pacific	17.8 ± 7.6		Steinberg et al. (1994)
Cyclopoid copepods	Coastal Pacific	68.7 ± 20.0		Steinberg et al. (1994)
Harpacticoid copepods	Coastal Pacific	23.2 ± 11.3		Steinberg et al. (1994)
<b>Protozoans</b>				
Flagellates	NW Atlantic	na	17–520	Caron et al. (1986)
Flagellates	NE Atlantic	$1.4 \times 10^3$ – $5.9 \times 10^3$	$1.3 \times 10^3$ – $3.3 \times 10^3$	Turley & Mackie (1994)
Choanoflagellates	Northern Adriatic Sea	$0.2 \times 10^5$ – $2.1 \times 10^6$ <sup>a</sup>	2–253	Herndl (1988)
Monads	Northern Adriatic Sea	$0.2 \times 10^5$ – $1.2 \times 10^6$ <sup>a</sup>	1–290	Herndl (1988)
Ciliates	Monterey Bay	0–130	$5$ – $8.6 \times 10^3$	Silver et al. (1978)
Ciliates	Monterey Bay	$1.5 \times 10^3$ – $1.7 \times 10^4$	30	Silver et al. (1998)
Sarcodines	Monterey Bay	220–230	220–380	Silver et al. (1998)
Flagellates	Elbe estuary	2–25	na	Zimmermann-Timm et al. (1998)
Ciliates	Elbe estuary	1–12	na	Zimmermann-Timm et al. (1998)
Amoebae	Elbe estuary	1–211	na	Zimmermann-Timm et al. (1998)

Table 2 (continued)

Organism	Location	Cells (ind.) agg <sup>-1</sup>	EF	Source
<b>Bacteria</b>				
	Southern California Bight	$1.3 \times 10^6$ – $1.7 \times 10^6$	30–37	Allredge et al. (1986)
	NW Atlantic	$1.8 \times 10^6$ – $2.8 \times 10^6$	20–55	Allredge et al. (1986)
	Monterey Bay	$4.2 \times 10^5$ – $6.7 \times 10^6$	83–300	Allredge et al. (1986)
	Monterey Bay	$3.2 \times 10^8$ – $5.4 \times 10^8$	0.6–16	Silver et al. (1998)
	NE Atlantic	$2.5 \times 10^5$ – $4.3 \times 10^5$	400–5,700	Turley & Mackie (1994)
	Southern California Bight	$0.1 \times 10^6$ – $1.7 \times 10^6$	na	Simon et al. (1990)
	Northern Adriatic Sea	$1.3 \times 10^6$ – $1.1 \times 10^8$ <sup>a</sup>	2–133	Herndl (1988)
	Lake Constance	$1.2 \times 10^6$ – $2.3 \times 10^7$	35–220	Grossart & Simon (1993)
	Elbe estuary	$5.0 \times 10^6$ – $5.0 \times 10^7$	na	Zimmermann-Timm et al. (1998)

<sup>a</sup>per ml agg; <sup>b</sup>% on agg

Table 3. Dyes for staining specific biopolymeric components on aggregates. DTAF: 5-(4,6-dichlorotriazinyl)aminofluorescein; FITC: fluorescein isothiocyanate; TRITC: tetramethyl rhodamine isothiocyanate

Dye	Specificity	Source
Alcian Blue	Acidic polysaccharides	Allredge et al. (1993), Logan et al. (1994)
Coomassie Blue	Proteins	Long & Azam (1996)
Toluidine Blue	Proteoglycans, acidic polysaccharides, polymers	Chayen et al. (1973)
DTAF	Polysaccharides	Schumann & Rentsch (1998)
FITC-/TRITC-labeled lectins	Specific bands of polysaccharides such as:	
Glycine max	<i>N</i> -acetyl-D-galactosamine	Grossart (1999)
Bandeiraea s. II	<i>N</i> -acetyl-D-glucosamine	Grossart (1999)
Wheat germ	<i>N</i> -acetyl-D-glucosamine, <i>N</i> -acetylneuraminic acid	Grossart (1999)
Concavaline A	Methyl $\alpha$ -D-mannopyranoside D (+) mannose	Grossart (1999), Neu (2000)
Abrin	D (+) galactose	Neu (2000)
Lentil	$\alpha$ -methyl-D-mannopyranoside D (+) mannose	Neu (2000)
Limulin	<i>N</i> -acetylneuraminic acid D-glucuronic acid	Neu (2000)
Lotus	L (-) fucose <i>N</i> -acetyl-D-glucosamine	Neu (2000)

tive function of the aggregate size except for larvacean houses, which are not formed by aggregation (Allredge & Silver 1988, Grossart & Simon 1993, Müller-Niklas et al. 1994, Allredge 1998, Grossart & Ploug 2000). Depending on the abundance, macroaggregates can contribute between <4 and ~30% to total POC in pelagic marine and lacustrine systems (Allredge & Silver 1988, Riebesell 1991a, Allredge et al. 1998, Grossart & Simon 1998b) but presumably much higher proportions in estuaries. Respective data, however, are not yet available.

The dominant biopolymers of macroaggregates are carbohydrates, proteins (or particulate combined amino acids [PCAA]) and lipids, constituting the majority of the aggregate-bound POM and also of the phytoplankton as the primary source of pelagic macroaggregates (Allredge 1979). Surprisingly little information, however, is available about the biochemical composition of macroaggregates of different origin. PCAA was 0.20 to 1.13  $\mu\text{g}$  per aggregate in selected marine snow samples from surface waters composed of larvacean houses and diatoms (Smith et al. 1992) and 0.14 to 0.47  $\mu\text{g}$  per aggregate in lake snow samples

(Grossart & Simon 1998b). These values constituted 8 to 51% of the aggregate-bound POM, which were higher proportions than of total POM (Grossart & Simon 1998b). This notion indicates that macroaggregates may be of higher nutritive value as food for particle feeders than total POM, in particular when algae with a low C:N ratio constitute only minor proportions of total POM. In fact, macroaggregates are a food source for various animals (see below). The microscopic and chemical analysis of coastal sedimenting material and of riverine macroaggregates in the Elbe estuary, Germany, indicated that substantial amounts of their POC consisted of macrophyte debris and lignin fairly refractory to microbial decomposition, except during phytoplankton blooms (Cowie & Hedges 1992, Zimmermann-Timm et al. 1998).

The composition of aggregates often is examined qualitatively by brightfield or epifluorescence microscopy after staining with dyes specific for certain biochemical compounds (Table 3). The recent and ongoing introduction of highly intense fluorescent dyes such as 5-(4, 6-dichlorotriazinyl)aminofluorescein (DTAF) and fluorescent lectins to detect specific poly-

saccharides will further help elucidate more details about the composition, fate and dynamics of organic micro- and macroaggregates (Schumann & Rentsch 1998, Grossart 1999, Neu 2000). Scanning (SEM) or transmission electron microscopy also has been used to examine the composition of micro- and macroaggregates (e.g. Paerl 1974, Leppard et al. 1979, Heissenberger et al. 1996, Grossart & Simon 1998b). It requires, however, fixing and drying of the specimen and thus may introduce artifacts. The introduction of confocal laser scanning microscopy (CLSM) provided new options to study the spatial organization of aggregates and, in combination with specific dyes and fluorescent lectins, allowed attractive and new important insights into the heterogeneous biochemical composition of aggregates (Caldwell et al. 1992, Manz et al. 1999, Neu 2000). In contrast to SEM it requires only little pretreatment of the specimen and allows it to be examined when fully hydrated. The combination of CLSM and fluorescent *in situ* hybridization with rRNA-targeted oligonucleotides further provides attractive means to study the spatial distribution of bacterial populations colonizing aggregates (Wagner et al. 1994, Manz et al. 1999, see Section 4.4.1).

### 3.4. Microaggregates

Microaggregates often appear quite similar to macroaggregates when examined microscopically even though their composition is not as diverse. Certain classes of microaggregates such as TEP and protein-containing particles (see Section 3.4.2), which may also include non-aggregated particles, are dominated by specific biopolymers. The abundance of microaggregates, however, is higher than that of macroaggregates by several orders of magnitude (Table 1) and they often exhibit somewhat different spatio-temporal distributions from macroaggregates, indicating that they are differently involved in the processes of POM cycling, i.e. formation and decomposition of macroaggregates. Their abundance and size are inversely related (Riebesell 1991a, Li & Logan 1995, Mostajir et al. 1995, Long & Azam 1996, Mari & Burd 1998, Worm & Søndergaard 1998b, Brachvogel et al. 2001). Due to their high abundance, microaggregates collide much more frequently with macroaggregates than the latter with each other. This implies that they are subject to removal by macroaggregates provided they have a high sticking efficiency or are captured efficiently. Knoll et al. (2001) examined the formation of microaggregates during a limnetic diatom bloom in Lake Constance and found that they are formed freshly from senescent diatoms and that their size increased from 50 to 80  $\mu\text{m}$  to  $>100 \mu\text{m}$  within 30 h.

These types of microaggregates, however, have only rarely been found *in situ*, presumably because of their high removal rate (Brachvogel et al. 2001). Azam et al. (1993), by examining the aggregation of marine diatoms, made similar observations of the initial formation of microaggregates and later of macroaggregates. Some types of microaggregates persist for quite a while and are not scavenged by macroaggregates, nor do they aggregate or sink, presumably because they have a low sticking efficiency and have a Reynolds number far below 1 (Drapeau & Dam 1995, Berman & Viner-Mozzini 2001, Brachvogel et al. 2001). In shallow turbid environments such as estuaries and tidally affected shallow seas with high hydrodynamic forcing, microaggregates exhibit the highest abundances and contain high amounts of resuspended inorganic particles (Fig. 2B). In the turbidity maximum of the Columbia River estuary on the Pacific NW coast of the USA, microaggregates in the size range 3 to 10  $\mu\text{m}$  were associated with 87% of total bacterial production, thus demonstrating a key role of this size fraction in the estuarine carbon cycling (Crump & Baross 2000). Hence, microaggregates can be involved very differently in the cycling of organic matter, depending on the given system, the load of suspended matter and the quality of the POM. Due to the small size it has not been possible to determine the individual amount of organic carbon or nitrogen of natural microaggregates. Recently, however, new stains specific for various biopolymers were introduced (Table 3), which enabled the determination of the biopolymer composition of organic aggregates and led to the discovery of new classes of microparticles outlined below.

#### 3.4.1. Transparent exopolymer particles

TEP were first described in 1993 from neritic waters off California (Alldredge et al. 1993). Thereafter they were found in many other environments including lakes and identified as a ubiquitous class of microparticles (Passow & Alldredge 1994, Passow et al. 1994, Logan et al. 1995, Schuster & Herndl 1995, Mari & Kiørboe 1996, Grossart et al. 1997, 1998, Kiørboe et al. 1998, Mari & Burd 1998, Worm & Søndergaard 1998b, Berman & Viner-Mozzini 2001, Brachvogel et al. 2001, Mari et al. 2001). TEP are stained bright blue by a simple technique with alcian blue (Alldredge et al. 1993, Logan et al. 1994). The abundance of TEP ranges from  $<10^2$  to  $>10^7 \text{ l}^{-1}$  (Table 1) and in general is positively correlated to phytoplankton biomass. The size varies between  $\sim 5$  and  $>100 \mu\text{m}$ , and size-class distributions can change quite rapidly in particular during aggregation events. Such aggregation events may clear the water column of TEP and appear to be one of the most

important mechanisms to reduce their abundance (Logan et al. 1995, Mari & Kiørboe 1996). Several field and experimental studies showed that TEP is produced by diatoms during the exponential growth or stationary phase (Kiørboe & Hansen 1993, Passow & Alldredge 1994, 1995, Passow et al. 1994, Schuster & Herndl 1995, Engel 2000). The amount of TEP produced is species specific and depends on the growth conditions. In some cases acidic polysaccharides do not detach from the surface of phytoplankton cells such that they directly stick together and form macroaggregates, and TEP is produced only in the course of the microbial break-down of the aggregates, possibly also from colloidal material (Kiørboe & Hansen 1993, Hansen et al. 1995, Passow & Alldredge 1995, Grossart et al. 1997, Mari & Burd 1998, Engel 2000). TEP associated with diatom aggregates obviously reduces their excess density and thus their sinking rate (Alldredge 1999, Engel & Schartau 1999). Thecate dinoflagellates also produce large amounts of mucopolysaccharides, but this material is not TEP because it does not stain with alcian blue (Alldredge et al. 1998). The C:N ratio of TEP, formed from diatom exudates by bubbling, has been determined to be close to 7, which is surprisingly low (Mari 1999). Because the experiments were carried out with non-axenic algae, TEP was possibly colonized by bacteria, thus reducing the C:N ratio. In a field study in the northwestern Mediterranean Sea, the C:N ratio of TEP produced from DOM by bubbling ranged between 6.5 and 70 (Mari et al. 2001). Mean values before and after a phytoplankton summer bloom were 11.6 as compared to 37.7 during the bloom.

Bacteria, in particular when attached to surfaces like biofilms or macroaggregates, produce large amounts of exopolysaccharides as well (Decho 1990, Costerton et al. 1995, Heissenberger et al. 1996, Stoderegger & Herndl 1998). When freely suspended, they were also reported to produce TEP and thus may be an important source of these microaggregates under non-bloom conditions of phytoplankton (Stoderegger & Herndl 1999). The general significance of heterotrophic bacteria in the production and cycling of TEP, however, is not yet known. Because all experimental studies on the formation of TEP by phytoplankton were carried out with non-axenic cultures and because under *in situ* conditions phytoplankton are always associated with bacteria, it is possible that bacteria are much more actively involved in the formation of TEP than we assume. Bacteria in the phycosphere may control the production of TEP by algae by reducing the availability of inorganic phosphate to algae, resulting in enhanced rates of algal polysaccharide secretion (Guerrini et al. 1998, Azam et al. 1999, Grossart 1999). To specifically examine the role of bacteria in forming

TEP, comparative experiments with axenic and non-axenic algae need be carried out. It has been suggested and inferred from TEP size spectra and coagulation models that TEP is formed of colloidal precursors, thus representing a significant sink of dissolved carbohydrates (Kiørboe & Hansen 1993, Passow & Alldredge 1994, Mopper et al. 1995, Mari & Burd 1998). Recently, it has been shown experimentally that TEP are formed from colloidal precursor material and that formation rates vary according to the growth conditions of phytoplankton blooms (Zhou et al. 1998, Passow 2000, Mari et al. 2001).

TEP has attracted much interest because it was identified as important in the aggregation of diatom blooms. Interestingly, a class of microparticles with the properties found in TEP was hypothesized and predicted to exist shortly before the discovery of TEP (Hill 1992). Coagulation theory often could not properly predict aggregation events at the end of phytoplankton blooms without postulating the existence of a class of microparticles with properties as discovered with TEP. In fact, TEP must be considered to be one of the most important biological factors controlling the aggregation of phytoplankton blooms, leading to the formation of macroaggregates. One case study in coastal waters found that a diatom-dominated phytoplankton community did not develop a bloom with a terminating sedimentation event because the concentration of TEP remained surprisingly low (Kiørboe et al. 1996). Instead, most algae were sinking packaged in fecal pellets or as intact cells attached to larvacean houses. The specific roles of diatoms and in particular of bacteria in forming, cycling and degrading TEP are not yet comprehensively understood and need further work, which certainly will reveal more interesting details.

#### 3.4.2. Coomassie blue-stained particles

Recently, a class of microparticles containing protein was discovered in the sea by staining water samples with Coomassie brilliant blue, a protein-specific dye (Long & Azam 1996). Their shape was quite variable and the size ranged from a few to  $>150\ \mu\text{m}$ . Their abundance ranged from  $\sim 10^6$  to  $3.5 \times 10^7\ \text{l}^{-1}$  (Table 1) and decreased vertically, suggesting a positive relationship with phytoplankton biomass. Interestingly, in a few comparisons, the abundance was systematically higher than that of TEP determined in parallel samples. Due to the incompatible staining of particles by alcian blue and Coomassie brilliant blue, a direct comparison with respect to their content of acidic polysaccharides and proteins is not possible. In the only other study available, Berman & Viner-Mozzini (2001)



investigated these protein-containing particles in Lake Kinneret. They found abundances ranging from  $\sim 10^4$  to  $\sim 10^7$  l<sup>-1</sup>, a close correlation of the abundance and area to the phytoplankton biomass and, in contrast to Long & Azam (1996), systematically lower numbers than of TEP. The general role, significance and fate of the protein-containing particles is not yet known. They may also be involved in the formation of macroaggregates but whether they exhibit a key function similar to that of TEP is at least questionable. Because protein is decomposed fairly rapidly, the microbial cycling of these particles may be higher than that of TEP (Grossart 1999). Further work is needed to better elucidate the origin and significance of these particles.

#### 3.4.3. DAPI-stained particles

A class of microparticles that are stained yellow by DAPI (4',6-diamidino-2-phenylindole), a DNA-specific fluorescent dye that stains double-stranded DNA blue and is normally used to stain and enumerate bacterial cells, has been described from Mediterranean waters (Mostajir et al. 1995). These particles occurred in the size range of 0.2 to 20  $\mu$ m, with abundances around  $2.2 \times 10^7$  l<sup>-1</sup> (Table 1), and with an exponentially decreasing size-class distribution. There was a trend of decreasing numbers with depth in the size classes <5  $\mu$ m but little variation with depth in the size classes 5 to 15  $\mu$ m. Even though some covariations with chlorophyll were found and some changes in the distribution with stratification and mixing, nothing is known about their origin, fate and role in aggregation.

Somehow, comparable microaggregates, stained blue or yellow with DAPI, were found in Lake Constance, Germany (Brachvogel et al. 2001). They were studied in the size range 5 to >60  $\mu$ m and exhibited abundances between  $3.1 \times 10^5$  and  $1.53 \times 10^6$  l<sup>-1</sup> (Table 1) with highest numbers in the smallest size class (8 to 20  $\mu$ m). They consisted predominantly of phytoplankton and zooplankton debris, diatom frustules and unidentifiable detrital components. The spatio-temporal patterns of these microaggregates differed from that of TEP, macroaggregates and diatoms, and exhibited some covariations with the biomass of chrysophytes and dinoflagellates. Brachvogel et al. (2001) suggest that they are the remains of detrital POM, either directly from senescent chrysophytes and dinoflagellates in surface waters, or from decomposed macroaggregates at greater depths. The conspicuous colonization by  $\beta$ -*Proteobacteria* and bacteria of the *Cytophaga/Flavobacteria* cluster but not by  $\alpha$ -*Proteobacteria* substantiated the fact that these microaggregates are rather refractory remains of detrital POM and produced at the depth where they were

found (see Section 4.4.1.2). Further studies have to better elucidate the origin, significance and role of these microaggregates, and how they are involved in the formation or break-down of macroaggregates.

## 4. COLONIZATION OF AGGREGATES BY HETEROTROPHIC ORGANISMS

From the previous sections it is obvious that the occurrence and formation of organic aggregates is rather variable and dynamic, as are their origin, size and composition. They undergo pronounced changes and successions, and interact in various ways with the surrounding organisms, particles and dissolved material. These dynamics and interactions exhibit differences depending on the given environment, e.g. the pelagic zone of lakes and the sea, estuaries or rivers. A closer examination shows that these changes, in particular after aggregates are formed, are greatly influenced by the activities of the aggregate-associated heterotrophic organisms and in particular microbes, and are reflected in the colonization patterns of the aggregates. Hence, the understanding of these changes is only as good as our knowledge of the activities and composition of the microbial communities involved.

### 4.1. Metazoans

Since macroaggregates were identified as important microenvironments for the cycling and decomposition of organic matter in aquatic ecosystems, many studies were carried out to study their colonization by various heterotrophic organisms. Besides algae (see above), a great variety of metazoans was found on marine, riverine and estuarine aggregates (Table 2). Colonization of marine snow aggregates by metazoans was first reported by Alldredge (1972) but only during the last 10 yr was the colonization of aggregates by metazoans studied more extensively (Shanks & Edmondson 1990, Bochdansky & Herndl 1992, Steinberg et al. 1994, 1997, Green & Dagg 1997, Shanks & Del Carmen 1997, Shanks & Walters 1997, Zimmermann-Timm et al. 1998, Kiørboe 2000). Most metazoans on aggregates are enriched in numbers as compared to the surrounding water, and aggregates in coastal waters harbor a substantial fraction of all metazoans (Table 2). So far, little is known about the general significance of metazoans for the cycling and decomposition of aggregates. An examination of the behavior of zooplankton in the presence of aggregates showed that most animals visited and fed on aggregates for only a few minutes, but the behavior of individual taxa exhibited

pronounced differences (Green & Dagg 1997, Shanks & Del Carmen 1997, Shanks & Walters 1997, Kiørboe 2000). The abundance of zooplankton on aggregates integrated over time implies much higher enrichment factors than presented in Table 2, in the order of  $10^2$  to  $10^4$ , 1 to 2 orders of magnitude higher than of protozoans and prokaryotes (Kiørboe 2000). These high enrichment factors cannot be explained by simple encounter rates or scavenging of zooplankton by sinking aggregates, and it has been proposed and shown that zooplankton detect sinking aggregates remotely by chemical clues due to the DOM plume leaking out (Kiørboe 2000, 2001, Kiørboe & Thygesen 2001). The swimming behavior of large zooplankton such as euphausiids can cause fragmentation of marine snow aggregates, whereas that of copepods and other smaller zooplankton cannot (Dilling & Alldredge 2000). Hence, the individual behavior of zooplankton in the surroundings of aggregates is important for their fate.

There is experimental evidence that zooplankton and juvenile fish and fish larvae actively feed on lake snow as well as marine snow and that the aggregate quality affects feeding and ingestion rates (Larson & Shanks 1996, Shanks & Walters 1996, 1997, Grossart et al. 1998, Dilling et al. 1998, Kamjunke & Mehner 2001, Kamjunke et al. 2002). The presence of TEP reduces ingestion of diatoms by euphausiids, but TEP are ingested as well as an alternative food source (Passow & Alldredge 2000). Coprophagy and coprophagy need also to be considered as processes by which zooplankton interact with, disrupt or ingest aggregates dominated by fecal material (Lampitt et al. 1990). Feeding by macrozooplankton and fish on macroaggregates is a direct link between the microbial loop and the higher trophic levels of the food web, because these animals reach bacteria associated with aggregates which they cannot ingest as freely suspended picoplankton cells. Because bacteria can constitute up to 50% of the protein of macroaggregates (Simon et al. 1990), they can be a major aggregate-associated food source for macrozooplankton even though phytoplankton algae are usually a preferred food source. A first estimate by Kiørboe (2000) indicates that decomposition rates of marine snow by associated invertebrate zooplankton are of similar magnitude to that by bacterial activity. Hence, zooplankton may play a significant role in reducing the export production from the euphotic zone due to fragmentation, consumption and mineralization of aggregates (Banse 1990, Lampitt et al. 1990, Dilling et al. 1998, Dilling & Alldredge 2000, Graham et al. 2000, Kiørboe 2000). Further studies are needed to better elucidate the role of metazoans for the decomposition of organic aggregates.

## 4.2. Fungi

Fungi are the primary decomposers of leaf litter and other macrophyte-derived detrital material in streams and rivers (e.g. Bärlocher 1992), and in the marine environment (e.g. Fell & Newell 1998). Therefore, they presumably also colonize micro- and macroaggregates in these ecosystems rich in macrophyte-derived organic matter. However, to our knowledge, nothing is known about the colonization of aggregates by fungi, which is a great gap in our understanding of the microbial ecology and decomposition of aggregates.

## 4.3. Protozoans

Protozoans such as heterotrophic flagellates, ciliates, sarcodines and amoebae have been found to colonize micro- and macroaggregates, and are highly enriched as compared to the surrounding water (Table 2). The protozoan community on aggregates exhibits differences from that in the surrounding water, because of the strikingly different environmental conditions, which obviously select more for benthic and biofilm-associated forms (Silver et al. 1978, Caron 1991, Rogerson & Laybourn-Parry 1992, Artolozaga et al. 1997, Zimmermann-Timm et al. 1998). Occasionally, nearly all flagellates and ciliates are found on aggregates and very few freely suspended in the water (Silver et al. 1978). It has been hypothesized that particles may be instrumental for the growth and survival of some species of flagellates and ciliates in the pelagic environment by grazing on particle-associated bacteria (Caron et al. 1982, 1986). Protozoans feed on aggregate-associated bacteria and algae. It has been shown that feeding by *Noctiluca scintillans* on diatom aggregates can be so intense that it prevents loss of aggregates due to sinking (Tiselius & Kiørboe 1998). Several studies showed that the population dynamics of protozoans follow that of bacteria and that they are important in regenerating ammonium and primary amines, which are released into the surrounding water (Pomeroy et al. 1984, Biddanda & Pomeroy 1988, Gotschalk & Alldredge 1989, Artolozaga et al. 1997, Grossart & Simon 1998a, Grossart & Ploug 2001). Hence, an important function appears to be remineralization of the aggregate-associated nitrogen and phosphorus, predominantly by grazing of bacteria. Interactions between protozoans and bacteria on aggregates, however, are still poorly studied. Recently, Ploug & Grossart (2000) showed that bacterial growth on aggregates can be balanced by bacterivory of protozoans. A further indication of intense bacterial grazing by protozoans is the occurrence of high abundances of aggregate-associated filamentous bacteria, which are known to be a

phenotypic response to protozoan grazing (Jürgens & Güde 1994, Grossart & Simon 1998b, Grossart & Ploug 2000).

#### 4.4. Prokaryotes

Prokaryotes, predominantly *Bacteria*, have been found on nearly all types of aggregates studied so far, including marine, lacustrine, riverine, and estuarine macro- and microaggregates. Microaggregates without any bacterial colonization were also reported (Berger et al. 1996, Long & Azam 1996, Zimmermann 1997, Brachvogel et al. 2001). Inorganic components such as calcite and clay particles are not usually colonized. The bacterial cells usually are not uniformly distributed on aggregates but often form microcolonies and occur in filamentous structures (e.g. Logan et al. 1994, Grossart & Simon 1998b, Grossart & Ploug 2000). The number of bacteria per aggregate is positively correlated to the aggregate size, ranging between <100 per aggregate on the smallest microaggregates (Passow & Alldredge 1994, Berger et al. 1996, Brachvogel et al. 2001, Knoll et al. 2001) and  $>10^6$  bacteria per aggregate on macroaggregates of a few to >10 mm, respectively (Table 2). Bacteria on aggregates are enriched as compared to the abundances of free-living bacteria in the surrounding water (Table 2). Enrichment factors decrease with increasing aggregate size (e.g. Alldredge et al. 1986, Davoll & Silver 1986, Herndl 1988, Alldredge & Gotschalk 1990, Turley & Mackie 1994, Ploug et al. 1999, Kjørboe 2000, Ploug & Grossart 2000). The bacterial abundance on aggregates is not proportional to the aggregate volume (Alldredge & Gotschalk 1990, Ploug et al. 1999, Ploug & Grossart 2000), and the number of bacteria per unit surface area is inversely related to the aggregate size, at least partially due to the fractal nature of the aggregates. Grossart & Simon (1993, 1998b) found that in Lake Constance the total number of aggregate-associated bacteria increased with depth, due to an increasing aggregate size. Grossart & Simon (1998b) and Herndl (1988) described a morphological succession of the bacterial community on aggregates with increasing age, indicating that aggregates undergo different decompositional stages.

The relative proportion of aggregate-associated bacteria to total bacterial numbers varies greatly, depending mainly on the abundance of aggregates. In most pelagic environments they constitute <10% and often <5% of total bacterial numbers (e.g. Alldredge & Gotschalk 1990, Grossart & Simon 1998b, Turley & Stutt 2000). In riverine and estuarine systems, however, they may constitute as much as 90% of total bacterial numbers and production (Bell & Albright

1981, Bent & Goulder 1981, Zimmermann & Kausch 1996, Zimmermann 1997, Crump & Baross 2000). In most environments, even with high amounts of suspended matter, the relative proportion of particle-associated bacteria is lower than that of free-living bacteria (Kirchman 1993, Berger et al. 1996, Crump & Baross 1996, Crump et al. 1998). Interestingly, it has been estimated that the number of bacteria colonizing TEP in pelagic marine environments equals 5 to 40% and in some cases 60% of the total bacterial numbers, suggesting that if TEP are not visualized by alcian blue many of these bacteria are enumerated as being free-living (Alldredge et al. 1993, Passow & Alldredge 1994, Mari & Kjørboe 1996, Worm & Søndergaard 1998b).

Bacteria on aggregates are often bigger than free-living bacteria in the surrounding water, ranging between 0.01 and 1.0  $\mu\text{m}^3$  or 30 to >100 fg per cell (Alldredge et al. 1986, Simon 1987, Herndl 1988, Alldredge & Gotschalk 1990, Simon et al. 1990). This is presumably due to the more favorable nutritive conditions than those in the surrounding water.

##### 4.4.1. Structural composition of aggregate-associated bacterial communities

A very important question to better understand the specific roles and functions of bacterial communities on aggregates concerns their structural composition. Due to the application of various molecular techniques, including the establishment of procaryotic clone libraries on the basis of sequencing the polymerase chain reaction (PCR)-amplified 16S rRNA gene or fragments of it, and fluorescence *in situ* hybridization with rRNA-targeted oligonucleotide probes of various specificity (FISH), fairly detailed and important information on the community structure of aggregate-associated bacterial communities has become available.

It must be kept in mind, though, that all the information about the structure of bacterial communities based on phylogenetic relationships does not reveal any trait of their physiological properties, except for a few phylogenetic clusters, such as the sulfate reducers (subcluster of  $\delta$ -*Proteobacteria*) or the ammonium oxidizers (subcluster of  $\beta$ -*Proteobacteria*). This drawback calls for a combination of phylogenetic and physiological analyses of bacterial communities. Due to very recent and ongoing methodological developments, the tools have become more and more available to perform such combined studies. Applying probes for specific aggregate-associated populations or physiological groups such as sulfate-reducing bacteria (SRB) (Grossart & Ploug 2000), methanogens or ammonium oxidizers, and combining them with substrate-specific fluores-

cent dyes and CLSM, or combining them with microautoradiography (Lee et al. 1999, Ouverney & Fuhrman 1999, Cottrell & Kirchman 2000) will result in new and exciting insights into the composition, the physiological activity and the spatial organization of aggregate-associated microbial communities.

#### 4.4.1.1. Marine environments

On the basis of terminal restriction fragment length polymorphism or fully or partially sequenced clone libraries of the 16S rRNA gene, the diversity of marine snow- and particle-associated bacterial communities as compared to free-living bacterial communities in the surrounding water has been studied in various marine environments and an estuary. A high diversity was found on marine snow with dominant phylotypes of the *Cytophaga/Flavobacteria* cluster, and  $\alpha$ - and  $\gamma$ -*Proteobacteria* (DeLong et al. 1993, Rath et al. 1998, Moeseneder et al. 2001). In contrast to these results, Acinas et al. (1999), in western Mediterranean waters, found only a low diversity in particle-associated bacterial communities with phylotypes belonging to the  $\gamma$ -*Proteobacteria*, and a higher diversity in free-living bacterial communities with phylotypes also from other phylogenetic lineages. In the same marine environment, Phillips et al. (1999) examined the composition of the ammonium-oxidizing bacterial community. They found that all particle-associated ammonium oxidizers belonged to the *Nitrosomonas* cluster whereas free-living ammonium oxidizers belonged to the *Nitrospira* cluster. Hence, this notion demonstrates that the same chemolithotrophic metabolism is performed by genetically distinctly different clusters in each habitat, obviously reflecting the different adaptational properties of the members of these clusters to the given environmental conditions. Bidle & Azam (2001) analyzed the bacterial community on aggregates of diatom detritus by denaturing gradient gel electrophoresis of PCR-amplified 16S rDNA fragments and found distinct temporal colonization patterns with a dominance of  $\gamma$ -*Proteobacteria* and *Sphingobacteria/Flavobacteria*.

The results mentioned above only give information on the qualitative composition of bacterial communities but no quantita-

tive data. Quantitative information is provided by analysis with FISH (Amann et al. 1995). Following this approach, members of the *Cytophaga/Flavobacteria* cluster were found to constitute around 30% of the DAPI-stainable cells on marine snow in the Southern California Bight (Ploug et al. 1999). Bacteria of this cluster, together with  $\gamma$ -*Proteobacteria*, were also dominant on naturally derived marine snow aggregates of various age in the polar frontal zone of the Southern Ocean, whereas  $\alpha$ -*Proteobacteria* constituted much lower proportions (Fig. 3). Only in 1 case were  $\alpha$ -*Proteobacteria* more important than  $\gamma$ -*Proteobacteria*, but cells of the *Cytophaga/Flavobacteria* cluster could not be enumerated in this and the other experiments in which no numbers of this cluster are shown. Bacteria of




Fig. 3. Composition of the bacterial community on macroaggregates at the Polar Front in the Southern Ocean. Upper panel: *Bacteria*; lower panel:  $\alpha$ -*Proteobacteria*,  $\gamma$ -*Proteobacteria*, *Cytophaga/Flavobacteria*. Hybridization with fluorescent oligonucleotides specific for *Bacteria* (EUB338),  $\alpha$ -*Proteobacteria* (ALF1b),  $\gamma$ -*Proteobacteria* (GAM42a) and *Cytophaga/Flavobacteria* (CF319) was done as described by Weiss et al. (1996). On the 5 samples from the left, numbers of CF319-labelled cells could not be enumerated. The macroaggregates were formed in rolling tanks of natural samples from various locations and depths and incubated over time under ambient light and temperature conditions. In one experiment aggregates were formed from *Thalassiothrix* sp. (Thal), which were collected by a net tow. Experiments were carried out in December 1995 and January 1996 on board RV 'Polarstern'. DAPI: 4',6-diamidino-2-phenylindole



all 3 phylogenetic lineages together constituted the great majority of the bacterial community and constituted >80% of the DAPI-stainable cells on the aggregates on which cells of the *Cytophaga/Flavobacteria* cluster were enumerated. The bacterial community on marine snow in the polar frontal zone was quite different from that in the surrounding water, which was largely dominated by cells of the *Cytophaga/Flavobacteria* cluster and comprised only very low proportions of *Proteobacteria* (Simon et al. 1999). A conclusion from the data so far available is that bacteria of the *Cytophaga/Flavobacteria* cluster, together with  $\gamma$ -*Proteobacteria*, appear to be dominant colonizers of marine snow aggregates. Bacteria of the former cluster are known to be able to degrade polymers including complex polysaccharides, which are important structural components of macroaggregates (see above, Reichenbach 1992).

#### 4.4.1.2. Lakes

The structure of bacterial communities on lake snow aggregates is strikingly different from that on marine snow even though the functional properties of the bacterial communities on both types of aggregates are rather similar (see below). Detailed studies in Lake Constance revealed that the bacterial communities on natural lake snow in the upper 25 m are dominated by  $\beta$ -*Proteobacteria*, which constitute 20 to 60% of the DAPI-stainable cells (Weiss et al. 1996, Grossart & Simon 1998b, Schweitzer et al. 2001). Applying 3 probes highly specific for the  $\beta$ -*Proteobacteria* *Duganella zoogloeoides* (formerly *Zoogloea ramigera*), *Acidovorax facilis* and *Hydrogenophaga palleroni*, Schweitzer et al. (2001) detected between 42 and 70% of all  $\beta$ -*Proteobacteria*.  $\alpha$ -*Proteobacteria* were also abundant on natural lake snow of an age of <2 d but usually did not exceed the abundance of  $\beta$ -*Proteobacteria*. By applying 2 probes highly specific for clusters of  $\alpha$ -*Proteobacteria* closely related to *Sphingomonas* spp. and *Brevundimonas diminuta* 16 to 60% of all  $\alpha$ -*Proteobacteria* were detected. These findings demonstrate that the aggregate-associated bacterial community in Lake Constance was dominated by surprisingly few species, even though nearly 200 clones were detected on aggregates (Huber 1997). This may reflect the close adaptation of a specialized bacterial community to the unique environmental conditions on aggregates. These specific populations already existed on microaggregates of an age of a few hours, which suggests that the seeding bacteria were already present on senescent algae prior to and during aggregation (Knoll et al. 2001). In support of this assumption, bac-

teria of these specific populations were never detected among the free-living bacteria (W. Zwisler & M. Simon unpubl. results). These seeding populations obviously include non-motile and gliding bacteria such as *D. zoogloeoides* (Dugan et al. 1992) and *Cytophaga* spp. (Reichenbach 1992), which presumably cannot actively approach the aggregates by chemotactic behavior such as many other motile bacteria (see below). The bacteria on aggregates detected by the highly specific probes or their close relatives are well known to colonize activated sludge flocs and thus indicate conspicuous similarities between these microenvironments (Wagner et al. 1993, 1994).  $\gamma$ -*Proteobacteria* were of no particular significance on lake snow in Lake Constance except on zooplankton debris and on phytodetrital aggregates composed of *Dinobryon* spp. (Brachvogel et al. 2001). Usually, they constituted <15% and often <5% of the DAPI-stainable cells. On aggregates of an age of >4 d, occurring below 30 m and on the sediment surface at 110 m, the relative proportion of  $\beta$ -*Proteobacteria* increased even more, but bacteria of the *Cytophaga/Flavobacteria* cluster occurred to increasing proportions of up to 15% (Schweitzer et al. 2001). These colonization patterns suggest that on the sinking aggregates the labile organic matter became increasingly decomposed, whereas the more refractory material resisted degradation. Interestingly, cells of the *Cytophaga/Flavobacteria* cluster always constituted higher proportions among the free-living bacterial communities (Zwisler & Simon 2002). Brachvogel et al. (2001) examined the bacterial colonization of microaggregates stained by DAPI (see above) and found that  $\beta$ -*Proteobacteria* and cells of the *Cytophaga/Flavobacteria* cluster dominated on these types of aggregates, constituting together 49 to 93% of *Bacteria*, which comprised 44 to 55% of the DAPI-stainable cells.  $\alpha$ -*Proteobacteria* were not detected at all on microaggregates. In line with the abovementioned results, these colonization patterns indicate that DAPI-stainable microaggregates were late decompositional stages of phytodetrital aggregates.

The examination of naturally derived macroaggregates by FISH revealed that in Lake Kinneret, Israel, the composition of the aggregate-associated bacterial community was rather similar to that in Lake Constance (B. Schweitzer & M. Simon unpubl. results).  $\beta$ -*Proteobacteria* and *Cytophaga/Flavobacteria* dominated as well, and *Duganella zoogloeoides* and *Acidovorax facilis* constituted substantial proportions of this *Proteobacteria* subclass. Hence, these colonization patterns appear to be a general pattern on lake snow aggregates.

Schweitzer et al. (2001) and Knoll et al. (2001) experimentally examined the colonization of naturally



derived lake snow aggregates and found a succession from an initially rather equal coexistence of  $\alpha$ - and  $\beta$ -*Proteobacteria* to a dominance of the latter within 2 d and decreasing proportions of  $\alpha$ -*Proteobacteria*. Concurrently, there was a transition from a consumption of dissolved amino acids by aggregate-associated bacteria to a release into the surrounding water (Schweitzer et al. 2001). After 2 to 3 d, bacteria of the *Cytophaga/Flavobacteria* cluster occurred on the aggregates with increasing proportions of up to 20%. These colonization patterns are similar to those found in the vertical distribution of bacterial communities on natural lake snow (see above) and reflect the decomposition process of the aggregates during sinking.

#### 4.4.1.3. Rivers and estuaries

In a study analyzing the bacterial community on aggregates in the River Elbe, Germany (using FISH), Böckelmann et al. (2000) showed that it was largely dominated by  $\beta$ -*Proteobacteria* constituting ~50% of total DAPI-stainable cells.  $\alpha$ - and  $\gamma$ -*Proteobacteria* were of lower significance and constituted 5 to 25% and 15 to 30%, respectively. Cells of the *Cytophaga/Flavobacteria* cluster made up not more than 15% of the DAPI-stainable cells from summer to winter but 40% in spring. Planktomycetes and SRB were also detected but usually constituted <10% except in fall, when the SRB fraction reached 18%. Using group-specific oligonucleotide probes, we examined the composition of the bacterial community on macroaggregates in the salinity gradient of the Elbe estuary in May and October (Fig. 4). In the limnetic section,  $\beta$ -*Proteobacteria* and cells of the *Cytophaga/Flavobacteria* cluster largely dominated, constituting 20 to 40% and 25 to 36% of the DAPI-stainable cells, respectively, in May, and 18 to 45% and 20%, respectively, in October.  $\alpha$ - and  $\gamma$ -*Proteobacteria* were of minor significance even though the former occasionally comprised 13% of the DAPI-stainable cells. These colonization patterns are rather similar to that of aggregate-associated bacteria found by Böckelmann et al. (2000) in a section of the River Elbe further upstream. They are also similar to patterns found in Lake Constance, even though the proportion of the *Cytophaga/Flavobacteria* cluster is greater, presumably because more refractory organic matter (macrophyte debris) is present in the River Elbe (Zimmermann-Timm et al. 1998).

The composition of the aggregate-associated bacterial community in the brackish and marine section of the Elbe estuary, however, was strikingly different from that in the limnetic section.  $\gamma$ -*Proteobacteria* largely dominated whereas  $\beta$ -*Proteobacteria* constituted not more than 12% (Fig. 4). Obviously,  $\beta$ -*Proteobacteria* cannot cope with the increase in salinity and do not grow any further, so that marine  $\gamma$ -*Proteobacteria* replace them. Further studies are needed to elucidate the closer phylogenetic affiliation of these marine aggregate-associated bacteria. In other studies the lack of  $\beta$ -*Proteobacteria* in the marine environment has also been described (Glöckner et al. 1999, Simon et al. 1999) but not yet their strong decrease in a salinity gradient toward the sea.




Fig. 4. Conductivity, total numbers of aggregate-associated bacteria, percentages of *Bacteria* (EUB338) and  $\alpha$ -(ALF1b),  $\beta$ -(BET42a) and  $\gamma$ -*Proteobacteria* (GAM42a), and *Cytophaga/Flavobacteria* cluster (CF319) at low and high tide in May 1997 along the salinity gradient of the Elbe estuary. Hybridization was done as described in Schweitzer et al. (2001)

In the Columbia River estuary, in the Pacific Northwest of the USA, a specific bacterial community on particles in the turbidity maximum dominated by phylotypes of the *Cytophaga/Flavobacteria* cluster and the  $\alpha$ -,  $\gamma$ - and  $\delta$ -*Proteobacteria* was found (Crump et al. 1999). The composition of this community was substantially different from that in the surrounding water but also from that on particles in the limnetic section of the estuary and the adjacent coastal ocean. Differences between the community composition of particle-associated and free-living bacteria were also found in Chesapeake Bay, USA (Bidle & Fletcher 1995).

Differences among the particle-associated bacterial communities in the Sacramento River, California, its estuarine turbidity maximum and the oligohaline reach of San Francisco Bay, were reported by Hollibaugh et al. (2000). These authors compared banding patterns of PCR-amplified 16S rDNA fragments separated by denaturing gradient gel electrophoresis. Differences between particle-associated and free-living bacterial communities were found to be minor, but they may have been overshadowed by an insufficient separation of the 2 fractions of the highly turbid water by filtration through 1.0  $\mu\text{m}$  filters.

## 5. MICROBIAL ACTIVITIES, SUBSTRATE TURNOVER AND AGGREGATE DECOMPOSITION

POM solubilization, substrate hydrolysis and uptake, biomass production, respiration and substrate release into the surrounding water are the major microbial, and predominantly bacterial, processes by which organic aggregates are decomposed (Fig. 5). Their relative significance varies among different types of aggregates, and ongoing research is continuously revealing important details of these processes to better understand the overall decomposition of aggregates, but also to identify very specific traits.

### 5.1. The aggregates' chemical microenvironment and solute exchange with the surrounding water

The application of microsensors and the development of a specially designed vertical flow system, in which individual macroaggregates are stabilized in the water by an upward directed flow opposing sinking, has made it possible to examine their chemical microenvironment, and thus mass transfer and microbial growth conditions in sinking macroaggregates (Ploug et al. 1997, Ploug & Jørgensen 1999, Ploug 2001). Comparisons of measured gradients of flow and solutes in the vicinity of aggregates in this flow system with those predicted from mass transfer theory revealed a close reproduction of the natural flow and diffusion field in the vicinity of sinking aggregates (Kiørboe et al. 2001, Ploug 2001). These authors demonstrated from theoretical considerations as well as from experiments that mass transfer to impermeable model particles and to aggregates is 4- to 20-fold facilitated as compared to non-moving aggregates, depending on aggregate size and sinking velocity, due to the exchange of water in the vicinity of sinking aggregates. The impact of flow on the exchange of solutes between an aggregate and the surrounding water may be even higher if the aggregates are porous and several millimeters to centimeters large. From theoretical considerations it has been shown that fluid flow within centimeter-large aggregates can increase the solute fluxes 2-fold from the surrounding water to cells in the interior of the aggregates, compared to that to free-living cells (Logan & Hunt 1987, Logan & Allredge 1989). Fluid flow velocities within macroaggregates have so far not been directly quantified because the techniques for such measurements are still under development. However, it was demonstrated recently that the bacterial production is up to 10-fold higher in sinking aggregates than in aggregates incubated under static conditions (see next section). Fluid flow around sinking aggregates thus is highly important for




Fig. 5. Loss processes and microbial decomposition pathways of macroscopic organic aggregates. POM: Particulate organic matter

the chemical microenvironment and growth conditions of the aggregate-associated bacteria and other microorganisms. The diffusion-limited oxygen supply and the oxidation potential of organic matter in millimeter-large aggregates in the pelagic environment is at least 1 order of magnitude higher than the diffusion-limited oxygen supply in sediments with similar volumetric respiration rates because of the large surface-to-volume ratio of suspended aggregates (Ploug & Jørgensen 1999).

## 5.2. Bacterial production

Rates of bacterial biomass production by the thymidine (Fuhrman & Azam 1980) and leucine methods (Kirchman et al. 1985, Simon & Azam 1989) have been measured on a variety of aggregates in marine as well as in limnetic systems (Table 4). Generally, they are positively correlated to the aggregate size (Alldredge & Gotschalk 1990, Grossart & Ploug 2000). Most of the cell-specific incorporation rates of thymidine or leucine or calculated bacterial growth rates were only slightly higher or in the same range as rates of free-living bacteria. Only in a few cases were significantly enhanced rates of aggregate-associated bacteria found (Alldredge & Gotschalk 1990, Simon et al. 1990, Smith et al. 1995, Grossart & Simon 1998b). Hence, estimated turnover times of aggregate-associated POC on the basis of bacterial biomass production are in the range of months to years and thus unrealistically long (Karl et al. 1988, Simon et al. 1990, Smith et al. 1992, Grossart & Simon 1993, 1998b). In combination with findings of

a high carbon demand of free-living bacteria in the mesopelagic zone (Cho & Azam 1988, Simon et al. 1992), they inspired studies to examine the solubilization and release of dissolved substrates from aggregates into the surrounding water. In fact, high solubilization rates and a net release of labile substrates into the surrounding water, exceeding greatly the carbon demand of aggregate-associated bacteria, were found in several studies that could explain rapid decomposition rates of aggregate-associated POM (Smith et al. 1992, Hoppe et al. 1993, Grossart & Simon 1998a,b).

Aggregate- or particle-associated bacterial biomass production as a proportion of total bacterial production in various ecosystems is highly variable and depends predominantly on the concentration of aggregates and suspended particles. In oligo- and mesotrophic pelagic marine and limnetic systems, it is usually <14% and only occasionally substantially higher, whereas in more eutrophic and estuarine systems with higher concentrations of aggregates it may exceed 30% (Table 4). In the Northern Adriatic Sea, Müller-Niklas et al. (1994) also found that aggregate-associated bacterial production exceeded that in the surrounding water.

Recent measurements of production rates by aggregate-associated bacteria led to a reconsideration of the relative significance of the aggregate-associated POC channeled into biomass production of the aggregate-associated bacteria and being released into the surrounding water. By incubating aggregates spiked with radiolabeled leucine and thymidine on a plankton wheel, where they were thus permanently suspended, bacterial production rates were up to 1 order of magni-

Table 4. Rates of bacterial biomass production (BP) and respiration on aggregates (agg) and aggregate-associated BP as percentage of total BP. nd: not determined

Location	Depth (m)	Bacterial production (ng C agg <sup>-1</sup> h <sup>-1</sup> ) (% of total BP)		Respiration (ng C agg <sup>-1</sup> h <sup>-1</sup> )	Source
Subtropical Atlantic	120–530	0.5–78.1	<0.39	nd	Alldredge & Youngbluth (1985)
NW Atlantic	7–15	0.23–13.8	2.3–26.0	nd	Alldredge et al. (1986)
Southern California Bight	7–15	0.68–8.03	1.3–13.4	nd	Alldredge et al. (1986)
Southern California Bight	7–15	<0.1–1.25	0.2–3.5	nd	Simon et al. (1990)
Southern California Bight	7–15	0.1–2.3	0.2–10.9	nd	Alldredge & Gotschalk (1990)
NE Atlantic	0–100	0.01–0.22	1.8–3.4	nd	Turley & Mackie (1994)
NW Mediterranean Sea	40–450	1.0–5.08	2.8–48.3	nd	Turley & Stutt (2000)
Lake Constance	5–25	<0.1–1.55	2–30	nd	Grossart & Simon (1993)
Lake Constance	5–25	<0.1–1.08	<1–13.9	nd	Grossart & Simon (1998b)
Frederiksborg Slottso	0.5	200–3770 <sup>a</sup>	9–70	nd	Worm & Søndergaard (1998a)
Columbia River estuary	0–13	400–2000 <sup>b</sup>	13–53	nd	Crump & Baross (1996)
Marine diatom aggregates	nd	6.3–187.5	nd	10.4–167	Ploug & Grossart (2000)
Riverine detrital aggregates	nd	<0.1–20.0	nd	1.0–89.6	Grossart & Ploug (2000)
Marine detrital aggregates	0–20	nd	nd	0.3–150	Ploug et al. (1999)
Marine diatom aggregates	0–10	nd	nd	3.6–120	Ploug (2001)

<sup>a</sup>>20 µm fraction (l<sup>-1</sup> h<sup>-1</sup>); <sup>b</sup>>10 µm fraction (l<sup>-1</sup> h<sup>-1</sup>)

tude higher than in non-suspended samples (Ploug & Grossart 1999, Grossart & Ploug 2000, Ploug & Grossart 2000). Traditionally, these measurements were carried out with pooled aggregates, which settled to the bottom of the test tubes (Alldredge et al. 1986, Alldredge & Gotschalk 1990, Simon et al. 1990, Smith et al. 1992, Grossart & Simon 1993, 1998b, Müller-Niklas et al. 1994). Examination of the oxygen gradient on the bottom of the test tube between the surrounding water and the center of the aggregate by a microelectrode revealed that a diffusive boundary layer had been established, which prevented a proper oxygen supply and rapid diffusion of the radiolabel to the inner parts of the aggregates (Ploug & Grossart 1999). Oxygen concentrations are up to 9-fold higher within 2 mm large sinking aggregates than in aggregates settling onto a solid surface, e.g. of an incubation vial or a sediment trap (Ploug & Jørgensen 1999). Hence, the unfavorable conditions lead to strongly reduced and thus biased production rates, which can be prevented by incubating the aggregates in test tubes rotating on a plankton wheel or a roller table. The production rates measured under these conditions may be somewhat higher than under *in situ* conditions because the oxygen supply *in situ* may not always be so favorable. They may also at least partially explain the notion that cell-specific thymidine incorporation rates are inversely related to the bacterial abundance normalized to microgram of aggregate (Alldredge & Gotschalk 1990). The conclusion from these experiments is that the heterotrophic bacteria on the aggregates utilize a substantially higher portion of the aggregate-associated POC than previously assumed, in particular amino-acid carbon (Grossart & Ploug 2001). This calls for a reexamination of the role of aggregate-associated bacteria for the cycling of aggregate-associated and total POC.

### 5.3. Respiration

In the studies in which carbon turnover by aggregate-associated bacteria has been determined, respiration has not been measured and bacterial growth efficiencies, if considered, were assumed according to published values of free-living bacteria or estimated indirectly (Smith et al. 1995, Grossart et al. 1998). Respiration of aggregate-associated microbes, though, can be determined from oxygen measured by a highly sensitive micro-Winkler technique (Parparov et al. 1998) or by oxygen microelectrodes (Alldredge & Cohen 1987, Ploug et al. 1997). The oxygen gradients around aggregates are proportional to the community respiration rates when mass transfer within aggregates occurs by molecular diffusion

(Ploug et al. 1997). Microelectrodes have been used to directly measure community respiration rates on marine snow from surface waters of the Southern California Bight, on riverine aggregates and on diatom aggregates (Table 4). In general, they were in the same range as, or slightly higher than, bacterial biomass production rates. On marine snow, respiration comprised  $75 \pm 21\%$  of the gross photosynthesis, even at light intensities  $>500 \mu\text{E m}^{-2} \text{s}^{-1}$ , thus emphasizing the dominance of heterotrophic processes on these aggregates. Generally, respiration was positively correlated with aggregate size but inversely correlated if respiration was normalized to aggregate volume, or per bacterium (Ploug et al. 1999, Ploug 2001). On riverine and naturally derived diatom aggregates, respiration rates were positively correlated to POC content and respiration was higher on aggregates of an age of 5 to 14 d as compared to 1 to 5 d (Grossart & Ploug 2000, Ploug & Grossart 2000). These authors calculated that the turnover of the aggregate-associated POC by respiratory losses was substantially higher than on the basis of just bacterial biomass production, thus indicating that respiration is an important pathway of aggregate decomposition, to which more attention should be addressed than previously.

### 5.4. Growth efficiency of aggregate-associated bacteria

When respiration on aggregates is predominantly due to bacteria, provided that protozoans do not particularly dominate, combined measurements of bacterial biomass production and respiration on the same aggregate allow an estimate of bacterial growth efficiencies of aggregate-associated bacteria. The few measurements carried out so far show that the growth efficiency is highly variable, with highest values of 0.4 to 0.5 in fresh aggregates and values decreasing to 0.05 in aggregates of an age of 14 d (Grossart & Ploug 2000). These data indicate that growth efficiencies of bacteria associated with aggregates rich in labile organic matter are at the higher end of values found in free-living bacterial communities and similar to those of bacteria growing on labile substrates (Del Giorgio & Cole 1998). In contrast, the few published growth efficiencies of aggregate-associated bacteria estimated indirectly by correlating bacterial biomass production with respiration (Parparov et al. 1998) and budgeting bacterial biomass production with primary production and carbon losses on aggregates (Smith et al. 1995) ranged between 0.09 and 0.17, and thus were at the lower end of values for free-living bacteria (Del Giorgio & Cole 1998).



### 5.5. Bacterial turnover of aggregate-associated organic matter

It has been well established for >15 yr that particle-associated bacteria exhibit higher cell-specific uptake rates of free amino acids and monosaccharides than free-living bacteria, which implies that the former are metabolically more active (for review see Kirchman & Ducklow 1987, Kirchman 1993). Detailed studies, however, on the turnover of labile substrates on aggregates, including hydrolysis, uptake and release, and determining total turnover rates of aggregate-associated carbon and nitrogen, only started about 10 yr ago, after Cho & Azam (1988) and Azam & Smith (1991) had postulated that microbial decomposition rates of aggregate-bound organic matter must be much higher than estimated on the basis of biomass production of aggregate-associated bacteria alone (see Section 5.2).

Several studies found that bacteria associated with marine and limnetic macroaggregates exhibit significantly higher potential ectoenzymatic hydrolysis rates per cell of aminopeptidase, phosphatase and glucosidases, implying that labile biopolymers on aggregates are hydrolyzed faster than in the surrounding water (Karner & Herndl 1992, Smith et al. 1992, 1995, Agis et al. 1998, Grossart & Simon 1998b, Unanue et al. 1998b, Worm & Søndergaard 1998a). Due to decomposition, solubilization and varying bacterial communities on aggregates, hydrolysis rates of these bacteria, though, are rather variable and, in particular on more decomposed and thus aged aggregates, are reduced and may be lower than those of their free-living counterparts (Müller-Niklas et al. 1994, Agis et al. 1998, Grossart & Simon 1998b, Unanue et al. 1998b, Grossart & Ploug 2001). Usually, per cell hydrolysis rates of aminopeptidase are higher than of  $\alpha$ - and  $\beta$ -glucosidase, but also decrease faster over time or depth, implying that protein is decomposed more rapidly than polysaccharides (Smith et al. 1995, Grossart & Simon 1998a, Simon et al. 2000). A preferential decomposition of organic nitrogen as compared to POC was observed by Grossart & Ploug (2001). These authors found a more rapid decrease of PCAA relative to total POC and enhanced uptake rates of amino acids relative to glucose on aggregates incubated in rolling tanks.

The rapid bacterial decomposition of PCAA and proteins has important implications for the decomposition of aggregates dominated by diatoms. Bidle & Azam (1999, 2001) demonstrated that dissolution of silica frustules in aggregates of diatom detritus takes place only when they are colonized by bacteria because they hydrolyze the protein matrix of the frustules. Individual bacterial species with different aminopeptidase activities exhibit different hydrolysis rates. Hence,

bacteria associated with diatom aggregates mediate a rapid recycling and enhance concentrations of silica within these aggregates (Brzezinski et al. 1997).

As a result of the high hydrolytic activities of aggregate-associated bacteria, and high metabolic activities of bacterial grazers, concentrations of DOC, dissolved free (DFAA) and combined amino acids (DCAA) and inorganic nutrients (phosphate, ammonium, nitrate, silicate) are significantly higher in the pore water of the aggregates than in the surrounding water (Table 5). Compared with DCAA, DFAA are relatively more enriched on aggregates than in the surrounding water (Table 5; Grossart & Simon 1998b), which is a further indication of the enhanced aminopeptidase activity of aggregate-associated bacteria. The high nutrient concentration in the pore water of aggregates, and the spatially tight coexistence of organisms within the aggregate matrix, enhanced nutrient recycling and uptake of the hydrolysis products by the aggregate-associated bacteria and algae and enhanced growth and biomass production rates (Gotschalk & Alldredge 1989, Alldredge & Gotschalk 1990, Ploug & Grossart 1999, Grossart & Ploug 2000, 2001). Besides recycling within aggregates, the highly enriched labile DOM and inorganic nutrients may also be partly released into the surrounding water. The finding that bacterial growth and production rates on aggregates often were not enhanced as compared to the surrounding water (see above) and that DCAA hydrolysis rates substantially exceeded the carbon and nitrogen demand of aggregate-associated bacteria led to the concept of an uncoupled hydrolysis and uptake of aggregate-associated organic matter by the associated bacteria and its subsequent predominant utilization by free-living bacteria in the surrounding water (Azam & Smith 1991). This concept was strongly supported by the measured net release of DFAA and DCAA from aggregates into the surrounding water (Smith et al. 1992, Grossart & Simon 1998a,b, Grossart & Ploug 2001, Schweitzer et al. 2001). Even though measured on various types of aggregates in marine and limnetic environments, DCAA release rates were relatively constant ( $0.32$  to  $2.25$  nmol  $\text{agg}^{-1} \text{h}^{-1}$ ). Please note that the respective data in Grossart & Simon (1998b) correctly range between  $0.45$  and  $2.25$  nmol  $\text{agg}^{-1} \text{h}^{-1}$  and thus are lower than those published in the original paper. Aggregates, however, do not always act as a source of dissolved amino acids but can also be a sink, as observed by Schweitzer et al. (2001) with fresh lake snow aggregates and by Grossart & Simon (1998a) with aged lake snow aggregates.

Recent model calculations and experiments, with a new approach simulating the release of DOM by sinking aggregates, indicate that the released compounds leave a plume with a strong spatial gradient behind the



Table 5. Enrichment factors (EF) of dissolved inorganic and organic nutrients on marine snow relative to the surrounding water. DOC: dissolved organic carbon; DFCHO: dissolved free monosaccharides; DTCHO: dissolved total carbohydrates; DFAA: dissolved free amino acids; DCAA: dissolved combined amino acids

Nutrient Location	Concentration (μM)		EF	Source
	Aggregates	Bulk water		
<b>Phosphate</b>				
Californian coastal Pacific	0.3–84	0.4–1.3	0.2–67	Shanks & Trent (1979)
Southern Californian Bight	0.04–21.7	0.01–0.4	<1–180	Allredge & Gotschalk (1990)
Northern Adriatic Sea	2.92 ± 5.78	0.1 ± 0.19	22	Kaltenböck & Herndl (1992)
Lake Constance	72–318	<0.1	720–3180	Grossart & Simon (1993)
<b>Nitrate</b>				
Californian coastal Pacific	0.4–308	0.6–12.6	11–24	Shanks & Trent (1979)
Southern Californian Bight	0.1–73	0.1–5.2	<1–108.5	Allredge & Gotschalk (1990)
Northern Adriatic Sea	3.84 ± 3.44	1.56 ± 1.91	2.5	Kaltenböck & Herndl (1992)
<b>Ammonium</b>				
Californian coastal Pacific	1–483	0.1–2.2	3–860	Shanks & Trent (1979)
Southern Californian Bight	0.2–25	0.1–0.4	5–55	Allredge & Gotschalk (1990)
Northern Adriatic Sea	4.03 ± 4.38	1.07 ± 1.35	1.9	Kaltenböck & Herndl (1992)
<b>Silicate</b>				
Monterey Bay	7.0–305	3.6–10.2	1.0–48	Brzezinski et al. (1997)
<b>DOC</b>				
Northern Adriatic Sea	1217.5	85	14.3	Herndl (1992)
Coastal NW-Pacific	742–11 700	108–150	16–78	Allredge (2000)
Southern Californian Bight	2430–2640	108–117	22	Allredge (2000)
<b>DFCHO</b>				
Northern Adriatic Sea	6.16	1.26–6.66	0.53–1.47	Herndl (1992)
<b>DTCHO</b>				
Northern Adriatic Sea	11.6–31.0	14.4	2.2	Herndl (1992)
<b>DFAA</b>				
Northern Adriatic Sea	0.49–0.54	0.25–0.56	0.96–1.96	Herndl (1992)
Northern Adriatic Sea	0.61–27.2	0.20	3.0–134	Kaltenböck & Herndl (1992)
Lake Constance	0.15–0.72	0.01–0.21	1.6–25.0	Grossart & Simon (1998b)
<b>DCAA</b>				
Lake Constance	2.3–49.9	2.0–9.5	0.9–11.1	Grossart & Simon (1998b)

aggregates, which can extend to several centimeters (Kjørboe 2001, Kjørboe et al. 2001). The volume with enhanced substrate concentration can be 10- to 100-fold larger than the volume of the aggregate itself. Sinking aggregates and their associated microbial processes thus create pronounced heterogeneities of nutrients in the bulk water, which may affect nutrient recycling and colonization processes by chemotactic microorganisms and remote chemosensory detection by meso- and macrozooplankton (Kjørboe 2001, Kjørboe & Jackson 2001, Kjørboe & Thygesen 2001, Kjørboe et al. 2001).

Even though production rates and respiratory carbon losses of aggregate-associated bacteria obviously are higher than assumed previously (see Section 5.2), the notion of a direct release of amino acids (see above), ammonium (Pomeroy et al. 1984) and DOM (Berman et al. 1999) strongly supports the validity of the concept that aggregates act as metabolically highly active bioreactors, leading to the decomposition and solubilization of the aggregate-bound POM by associated

microbes, but also by bacteria in the surrounding water, which decompose the released dissolved substrates. Recent experimental studies with laboratory-made aggregates examining the time course of aminopeptidase,  $\alpha$ - and  $\beta$ -glucosidase activities, and leucine and glucose uptake by aggregate-associated and free-living bacteria further support this concept (Agis et al. 1998, Unanue et al. 1998b). These authors found that ratios of aminopeptidase/bacterial production, aminopeptidase/leucine uptake and  $\beta$ -glucosidase/glucose uptake by free-living bacteria were initially rather low and increased later than those by aggregate-associated bacteria. Further, concentrations of DFAA in the surrounding water increased over time and covaried with the ratio of aminopeptidase/leucine uptake by aggregate-associated bacteria and thus implied intense DCAA hydrolysis.

Measurements of concentration dynamics of dissolved amino acids in the surrounding water of freshly formed lake snow aggregates, however, showed that a net release of amino acids usually occurs with aggre-

gates of an age of at least 2 d (Schweitzer et al. 2001). These authors observed that the release phase was usually preceded by a period in which aggregate-associated bacteria consumed amino acids from the surrounding water. Hence, it is obvious that fresh aggregates in the mixed layer can also act primarily as a sink for labile DOM such as amino acids, convert it rather effectively into bacterial biomass due to a high growth efficiency (see above) and transport it, as a fraction of total aggregate-associated POM, to deeper layers, where a substantial proportion is reconverted to DOM and serves to support the growth of the free-living bacterial community.

So far, only very few studies have been carried out to directly compare the relative significance of bacterial production, respiration and solubilization for the overall decomposition of aggregates. In one study, which simultaneously measured bacterial production, respiration, net release of amino acids and ectoenzymatic activity on individual diatom aggregates, bacterial production and mineralization were found to be the major decomposition pathways for carbon and nitrogen (Grossart & Ploug 2001). A direct comparison of respiration and solubilization of organic matter of lake snow aggregates of various origins showed that both processes were highly variable, such that in some cases respiration dominated and in others solubilization dominated (Berman et al. 1999). More studies are needed to better understand under which conditions either of the processes mentioned is favored or suppressed.

#### 5.6. Anoxic processes of aggregate-associated prokaryotic populations

Methanogens, and ammonium-oxidizing and nitrifying bacteria have been detected on and isolated from oceanic particulate matter, fecal pellets and zooplankton (Oremland 1979, Bianchi et al. 1992, Marty 1993, Phillips et al. 1999), and anoxic macroaggregates colonized by methanogens have been suggested as a source for the observed supersaturation of methane in the ocean (Karl & Tilbrook 1994). The production of CO<sub>2</sub> and ammonium within aggregates can promote growth of nitrifying bacteria, as has been observed in the mesopelagic zone in the North Pacific (Karl & Tilbrook 1984), and aggregates can be significant sites of ammonium oxidation by nitrifying bacteria in estuaries with high ambient ammonium concentrations (Kerner & Gramm 1995, Stehr et al. 1995, Schäfer & Harms 1996). The efficient mass transfer even to impermeable sinking aggregates and the high porosity of macroaggregates, however, imply that sinking aggregates are rarely anoxic in the natural environ-

ment. In non-sinking fecal pellets, i.e. not exposed to any water flow, anoxic microzones have been found by Alldredge & Cohen (1987). Concentration gradients of oxygen not reaching anoxia, however, do develop within sinking aggregates when microbial activities are high (see above). Even the diffusion-limited oxygen supply from the ambient water at air saturation can theoretically oxidize all aggregate-associated POC to CO<sub>2</sub> within 1 d, in 0.1 to 10 mm large sinking aggregates (Ploug et al. 1997). Such high metabolic rates are rarely observed in aggregates, and anoxic aggregates are therefore most likely to occur in the oxygen minimum zone in the ocean (Shanks & Reeder 1993, Ploug et al. 1997, 1999, Ploug 2001).

A conclusion from the mentioned observations is that anoxic processes may occur on certain types of macroaggregates under special environmental conditions but in general appear to be an ephemeral phenomenon and therefore presumably do not play a major role in the aggregate-associated decomposition of organic matter.

### 6. BACTERIA-BACTERIA AND BACTERIA-ALGAE INTERACTIONS

From the previous sections it is evident that aggregate-associated bacteria colonize highly specific microenvironments, which can be regarded as hot-spots of microbial activities and turnover of POM in various aquatic systems. An important question in this context concerns the way bacteria first colonize algae, or micro- or macroaggregates. As very specific bacterial populations develop on aggregates (see above), they must first sense their potential colonization site and be able to get into close contact with it. This process can only be successful if such bacteria are motile or if they already inhabit the precursor component particles. Recent studies on bacterial motility demonstrate that various marine bacteria have specific flagellin genes (Winstanley & Morgan 1997), and that natural assemblages of marine bacteria exhibit high-speed motility and large accelerations after enrichment with nutrients (Mitchell et al. 1995, Fenchel 2001, Grossart et al. 2001, Kiørboe & Jackson 2001). Both characteristics are prerequisites for the chemotactic behavior and active interactions between bacteria and particulate colonization sites such as phytoplankton cells and suspended aggregates.

Bacteria can be loosely or tightly associated with phytoplankton (ZoBell 1941, Sieburth 1968, Caldwell & Caldwell 1978), leading to a multitude of possible interactions between these organisms in aquatic ecosystems (see review by Cole 1982). Symbiotic interactions between phytoplankton and heterotrophic bacteria led

Bell & Mitchell (1972) to formulate the concept of the 'phycosphere': chemotactic bacteria benefit from extracellular polymers, mainly polysaccharides, which are released by the algae, whereas the algae benefit from the nutrients remineralized by the associated microbes (Golterman 1972), vitamins (Pringsheim 1912, Haines & Guillard 1974) and other growth factors (Paerl & Pinckney 1996). Bacteria in the phycosphere can be free-living, be directly attached to algal surfaces or live inside algal cells (Kochert & Olson 1970, Kodama et al. 1990, Rausch de Traubenberg & Lassus 1991). It has been shown that marine bacteria accumulate around phytoplankton-derived detritus, resulting in increased microbial activities in the 'detritosphere' (Biddanda & Pomeroy 1988). Recent theoretical models of chemotactic behavior of marine bacteria support the hypothesis that bacteria, clustering around phytoplankton or macroaggregates, greatly benefit from increased concentrations of nutrients in the close vicinity of otherwise nutrient-poor environments (Blackburn et al. 1997, 1998, Kiørboe & Jackson 2001). New approaches to studying the release of dissolved substrates, e.g. amino acids, from marine snow, show that sinking organic aggregates are followed by a plume of released DOM, which extends the aggregate length and enlarges its volume many times (Kiørboe et al. 2001). Such plumes of potential bacterial substrates, e.g. amino acids and oligopeptides, can be detected by chemotactic bacteria and lead to intense clustering of motile bacteria around these nutrient point sources (Kiørboe 2001, H. P. Grossart & F. Azam unpubl. results). Larger organisms such as copepods may sense these nutrient point sources as well (Kiørboe & Thygesen 2001).

Phytoplankton and heterotrophic bacteria, however, do not always benefit from each other (see review Cole 1982, Imai et al. 1993). It has been shown that heterotrophic bacteria can be repelled by exudates of the algae, e.g. antibiotics (Steemann-Nielsen 1955, Sieburth 1968), which has been interpreted as a mechanism by which living phytoplankton can prevent bacterial colonization of their cell surface. It has been further shown that a former symbiotic relationship between phytoplankton and heterotrophic bacteria can turn into a parasitic one when the alga becomes stressed, e.g. by missing vitamins (Grossart 1999). Different species of attached bacteria may also compete for space and nutrients on the aggregate surface, e.g. outcompeting each other by increased growth rates or the production of growth- or motility-inhibiting secondary metabolites (Kawano et al. 1997, Burchard & Sorongon 1998, Long & Azam 2001). Further studies on specific bacteria-bacteria and bacteria-algae interactions are needed and will lead to a more precise understanding of the microbial colonization and decomposition of aggregates.

## 7. VIRUSES

In the recent past, viruses were identified as an integral component of aquatic ecosystems, causing substantial proportions of the phytoplankton and bacterioplankton mortality (see review by Wommack & Colwell 2000). However, there is very little information available on the significance of virus infection on aggregate-associated bacteria and nothing on algal infection. Proctor & Fuhrman (1991) reported that in aggregated POM collected in sediment traps deployed in the northeastern Pacific, between 30 and 140 m depth, 0.7 to 3.7 % of the bacteria were visibly infected by viruses. In POM collected in a sediment trap deployed at 400 m, they did not find any infected bacteria. These proportions of infected bacteria are similar to values in free-living planktonic bacteria (Wommack & Colwell 2000). Hence, these observations indicate that virus infection of bacteria in marine snow aggregates is of similar significance to bacterial mortality as in free-living bacterial communities.

We examined numbers of bacteria and virus-like particles in lake snow aggregates formed in rolling tanks from Lake Constance water and incubated for up to 144 h (Fig. 6). The numbers of bacteria per aggregate increased over time in the experiments in May and August but not in June. Numbers of virus-like particles, counted by an epifluorescence standard method (Hennes & Suttle 1995), in the May experiment also continuously increased and covaried with bacterial numbers. In the June experiment virus-like particles decreased from initially high numbers until 23 h and covaried with bacterial numbers. In the experiment in August, numbers of virus-like particles peaked after 2 d but did not covary with bacterial numbers. Ratios of virus/bacteria generally followed dynamics of virus-like particles and ranged between 0.3 and 8.5 with a mean of  $4.7 \pm 2.3$ . In the experiments in May and August they were closely correlated to virus numbers ( $r^2 = 0.71$  and  $0.89$ ,  $p < 0.01$ ). These ratios are in the same range as values reported from studies on dynamics of viruses and free-living bacteria (Wommack & Colwell 2000) and imply that virus infection of bacteria and virus-induced bacterial mortality on aggregates is similar to that in free-living bacterial communities, confirming the results of Proctor & Fuhrman (1991) for marine snow aggregates. The rapid decreases we found suggest that viruses are readily decomposed in aggregates, possibly by the highly active bacterial aminopeptidases of the protein capsid (see above). Hence, large aggregates presumably are not a vehicle for free viruses to deep water layers and the bottom sediment. Only infected bacteria may bring viruses to these deep environments.

## 8. THE SIGNIFICANCE OF ORGANIC AGGREGATES TO THE FLUX AND CYCLING OF POM

From the presented information it is evident that organic aggregates and associated microorganisms are involved in many ways in the decomposition, cycling and flux of POM, DOM and inorganic nutrients in aquatic ecosystems. Major differences occur between pelagic marine and limnetic systems, with rather weak hydrodynamic forcing, and shallow and turbid marine, estuarine, riverine and lacustrine systems with strong and highly variable hydrodynamic forcing. The general aggregate-related microbial processes are rather similar in the various systems, even though the origin, size and composition of the aggregates and the major groups of microorganisms colonizing and decomposing them, and their quantitative significance for the overall POM cycling, may exhibit pronounced differences. In the following, and as a synopsis of this review, we want to highlight the major and most important features and the special significance of aggregate-related processes in the systems mentioned.

### 8.1. Pelagic marine and limnetic systems

A synopsis of formation and decomposition processes of marine and lake snow aggregates in pelagic systems with low hydrodynamic forcing is given in Fig. 7. Macroaggregates in these systems are rather large, 500  $\mu\text{m}$  to >10 mm, but in lakes they are smaller than in marine systems, presumably due to the lower bridging potential of divalent cations in the latter. De novo production can originate from various sources, including abandoned larvacean houses, pteropod webs and gelatinous phytoplankton such as *Phaeocystis* in marine systems and colony-forming cyanobacteria in lakes. Biologically and chemically mediated physical aggregation may include a large variety of component particles but phytoplankton often dominates. However, only a small proportion of the phytoplankton and primary production is usually in aggregated form.

The role of bacteria in aggregation still needs further studies, but bacteria may be of higher importance in producing TEP or controlling TEP production by diatoms than assumed, and may also directly control aggregation. The aggregate abundance in pelagic




Fig 6. Time courses of numbers of bacteria and virus-like particles (VLP), and the ratio of VLP/bacteria on aggregates formed from Lake Constance water in rolling tanks and incubated at ambient light and temperature conditions. Results of experiments in May 1996 (top), June 1995 (middle) and August 1996 (bottom) are shown. Note the different scales for bacteria, virus-like particles and time

systems is rather low (<130  $\text{l}^{-1}$ ) and hence aggregate-associated microbial processes appear to be of relatively low overall importance. However, it must be kept in mind that macroaggregates are surrounded by a phycosphere and a plume of nutrients and microbes, extending their size considerably and also affecting microbial and zooplankton-related processes in the surrounding water (Fig. 5; Azam & Long 2001, Kjørboe 2001). Because of this microheterogeneity and the total space that is occupied by aggregate-related processes, their overall significance presumably is much greater than assumed from traditional measurements and is still underestimated. The aggregate-associated microbial community exhibits pronounced differences from that in the surrounding water. On lake snow aggregates, bacteria are dominated by  $\alpha$ - and in particular by  $\beta$ -*Proteobacteria*, and cells of the *Cytophaga/Flavobacteria* cluster constitute increasing




Fig. 7. Formation and loss processes of macroscopic organic aggregates in pelagic ecosystems with low hydrodynamic forcing (modified from Alldredge & Silver 1988). The size of the arrows and letters reflects the relative significance of the respective pathway. For microbial decomposition pathways see Fig. 5. DIM: dissolved inorganic matter; DOM: dissolved organic matter; COM: colloidal organic matter




Fig. 8. Formation and loss processes of macroscopic and microscopic organic aggregates in shallow turbid ecosystems with strong hydrodynamic forcing. The size of the arrows and letters reflects the relative significance of the respective pathway. For microbial decomposition pathways see Fig. 5, definitions as in Fig. 7



proportions on aged aggregates with increasing fractions of refractory organic compounds. On marine snow,  $\gamma$ -*Proteobacteria* and cells of the *Cytophaga/Flavobacteria* cluster appear to be the major bacterial colonizers, but nothing is known about the temporal succession.

Even though, from a quantitative perspective, macroaggregate-associated decomposition processes appear to be of minor importance as compared to the total POM decomposition, macroaggregates have to be considered as hot-spots and thus of key importance for the whole pelagic ecosystem. Nutrients limiting the growth of phytoplankton and bacteria, such as inorganic nitrogen, phosphorus and silica, are highly enriched and recycled very rapidly in aggregates, thus also supplying free-living phyto- and bacterioplankton with regenerated nutrients (Fig. 5), and possibly controlling aggregation of phytoplankton (Fig. 7). Macroaggregates dominate the sinking flux of POM in pelagic ecosystems and thus are of key importance in supplying nutrients and energy to the pelagic and benthic biota in deep aphotic layers.

## 8.2. Shallow turbid marine, estuarine and limnetic systems

Aggregate formation in shallow turbid systems with high hydrodynamic forcing exhibits pronounced differences from pelagic systems with low hydrodynamic forcing (Fig. 8). Generally, the suspended matter load in these systems is much higher than in pelagic systems including also the proportion of primary production in aggregated form. De novo production originates from gelatinous phytoplankton, and pelagic and benthic fecal pellets, but also from benthic biofilms disrupted and resuspended by high shear and turbulence. Biologically and chemically mediated physical aggregation includes a large variety of component particles, not only from the water column but also resuspended from the sediment. In addition to those from the water column, benthic algae, debris of macroalgae and vascular plants, and high amounts of resuspended inorganic particles constitute major aggregate fractions. The dominant physical process causing aggregation of the component particles is collision by turbulent shear of rather high intensity, not allowing large aggregates to persist for longer periods, but disrupting them again such that microaggregates dominate. Differential settling is only important for short periods of time with low wind-induced shear, or at slack water in tidally affected systems. Resuspension of sediment components and intense restructuring of the aggregates because of the high hydrodynamic activities are major features of these systems.

Biological activities on biofilms and of the zoobenthos are also important for aggregate dynamics in shallow turbid systems including the nepheloid layer. The aggregates in these systems are colonized, consumed and repackaged by various metazooplankton and protozoans, including high proportions of benthic and meroplanktonic forms. The aggregate-associated bacterial community in shallow turbid limnetic systems is dominated by  $\beta$ -*Proteobacteria* and cells of the *Cytophaga/Flavobacteria* cluster, thus reflecting the dominance of rather refractory material in the aggregate-associated POM. In marine systems,  $\gamma$ -*Proteobacteria* and cells of the *Cytophaga/Flavobacteria* cluster obviously dominate, even though further studies are needed to confirm the existing preliminary results. Microbial decomposition processes on aggregates in general are similar to pelagic systems, but their overall significance with respect to total POM decomposition is much more important and may even be dominant. To better understand their relative importance, more studies examining these processes are needed. From the information presented in this review it is obvious that aggregate-associated microbial decomposition processes are highly significant in these systems and lead to solubilizing and mineralizing the POM, thus exporting its components as DOM and inorganic nutrients from estuaries and tidally affected shallow coastal areas to the coastal sea, or from the littoral to the pelagic zone of lakes. Aggregate consumption by the zoobenthos, meroplankton and fish appears to be highly significant, showing that this detrital and microbial food source supplies high amounts of energy and nutrients to higher trophic levels.

## 9. CONCLUSIONS

Today, the microbial ecology of organic aggregates in marine and limnetic systems appears to be rather well understood, and the major processes forming and decomposing aggregates have been identified and studied in detail. Further work, in particular on specific details with respect to the role of bacteria in forming aggregates and interacting with algae, will certainly improve our understanding of the microbial ecology of organic aggregates and their significance in cycling of nutrients and elements, and the flux of energy in aquatic ecosystems more so. Only very recently have aggregate-associated processes been included in food web and carbon cycling models of pelagic systems (Jackson 2001). The results of this first modeling assay suggest that aggregation of phytoplankton strongly affects the relative partitioning of zooplankton grazing and sedimentation of primary production. Such approaches need to be elaborated and tested much

more and to be applied to other aquatic ecosystems. The inclusion of organic aggregates in food web models and concepts is urgently needed for a more comprehensive understanding of these systems, and will give us a much better and more realistic understanding of ecosystem functioning in limnetic, as well as marine systems, from selected local systems to a global scale.

**Acknowledgements.** We are most grateful to the crew of RV 'Polarstern' and to our colleagues on board for their excellent cooperation during cruise ANT XIII/2. We would like to thank H. Zimmermann-Timm for her collaboration in the study of the Elbe estuary and providing us with conductivity data. Constructive suggestions on an earlier version of this review by A. L. Alldredge and 2 anonymous reviewers are gratefully acknowledged. This work was supported by grants from the Deutsche Forschungsgemeinschaft awarded to M.S. (Si 360/4-2 /4-3, Si 360/7-2), and from the Danish Natural Science Research Council (9801391) and the Carlsberg Foundation (990536/20-950 and -513) awarded to H.P.

#### LITERATURE CITED

- Acinas SG, Antón J, Rodríguez-Valera F (1999) Diversity of free-living and attached bacteria in offshore western Mediterranean waters as depicted by analysis of genes encoding 16S rRNA. *Appl Environ Microbiol* 65:514–522
- Agis M, Unanue M, Irriberry J, Herndl GJ (1998) Bacterial colonization and ectoenzymatic activity in phytoplankton-derived model particles. Part II. Cleavage and uptake of carbohydrates. *Microb Ecol* 36:66–74
- Aksnes DL, Wassmann P (1993) Modeling the significance of zooplankton grazing for export production. *Limnol Oceanogr* 38:978–985
- Alldredge AL (1972) Abandoned larvacean houses: a unique food in the pelagic environment. *Science* 17:885–887
- Alldredge AL (1979) The chemical composition of macroscopic aggregates in two neritic seas. *Limnol Oceanogr* 24:855–866
- Alldredge AL (1991) In situ collection and laboratory analysis of marine snow and large fecal pellets. In: Hurd DC, Spencer DW (eds) *Marine particles: analysis and characterization*. American Geophysical Union, Washington, DC, 43–46
- Alldredge AL (1992) Marine snow in ocean cycling. In: Nierenberg WA (ed) *Encyclopedia of earth system science*, Vol 3. Academic Press, San Diego, p 139–147
- Alldredge AL (1998) The carbon, nitrogen and mass content of marine snow as a function of aggregate size. *Deep-Sea Res* 45:529–541
- Alldredge AL (1999) The potential role of diatom exudates in forming nuisance mucilaginous scums. *Ann Ist Super Sanita* 35:397–400
- Alldredge AL (2000) Interstitial dissolved organic carbon (DOC) concentrations within sinking marine aggregates and their potential contribution to carbon flux. *Limnol Oceanogr* 45:1254–1253
- Alldredge AL, Cohen Y (1987) Can microscale chemical patches persist in the sea? Microelectrode study of marine snow, fecal pellets. *Science* 235:689–691
- Alldredge AL, Crocker KM (1995) Why do sinking mucilage aggregates accumulate in the water column? *Sci Total Environ* 165:15–22
- Alldredge AL, Gotschalk C (1988) In situ settling behavior of marine snow. *Limnol Oceanogr* 33:339–351
- Alldredge AL, Gotschalk C (1989) Direct observation of the mass flocculation of diatom blooms: characteristics, settling velocities, and formation of diatom aggregates. *Deep-Sea Res* 36:159–171
- Alldredge AL, Gotschalk C (1990) The relative contribution of marine snow of different origin to biological processes in coastal waters. *Cont Shelf Res* 10:41–58
- Alldredge AL, Madin LP (1982) Pelagic tunicates: unique herbivores in the marine plankton. *Bioscience* 32:655–661
- Alldredge AL, McGillivray PA (1991) The attachment probabilities of marine snow and their implications for particle coagulation in the ocean. *Deep-Sea Res* 38:431–443
- Alldredge AL, Silver ML (1988) Characteristics, dynamics and significance of marine snow. *Prog Oceanogr* 20:41–82
- Alldredge AL, Youngbluth MJ (1985) The significance of macroscopic aggregates (marine snow) as sites for heterotrophic bacterial production in the mesopelagic zone of the subtropical Atlantic. *Deep-Sea Res* 32:1445–1456
- Alldredge AL, Cole JJ, Caron DA (1986) Production of heterotrophic bacteria inhabiting macroscopic marine aggregates (marine snow) from surface waters. *Limnol Oceanogr* 31:68–78
- Alldredge AL, Gotschalk C, MacIntire S (1987) Evidence for sustained residence of macrocrustacean fecal pellets in surface waters of Southern California. *Deep-Sea Res* 34:1641–1652
- Alldredge AL, Granata TC, Gotschalk C, Dickey TD (1990) The physical strength of marine snow and its implications for particle disaggregation in the ocean. *Limnol Oceanogr* 35:1415–1428
- Alldredge AL, Passow U, Logan BE (1993) The abundance and significance of a class of large transparent organic particles in the ocean. *Deep Sea Res* 40:1131–1140
- Alldredge AL, Gotschalk C, Passow U, Riebesell U (1995) Mass aggregation of diatom blooms: insights from a mesocosm study. *Deep-Sea Res II* 42:9–27
- Alldredge AL, Passow U, Haddock HD (1998) The characteristics and transparent exopolymer particle (TEP) content of marine snow formed from thecate dinoflagellates. *J Plankton Res* 20:393–406
- Amann RI, Ludwig W, Schleifer KH (1995) Identification and in situ detection of individual microbial cells without cultivation. *Microbiol Rev* 59:143–169
- Artolozaga I, Santamaria E, Lopez A, Ayo B, Irriberry J (1997) Succession of bacterivorous protists on laboratory-made marine snow. *J Plankton Res* 19:1429–1440
- Asper VL (1987) Measuring the flux and sinking speed of marine snow aggregates. *Deep-Sea Res* 34:1–17
- Azam F (1998) Microbial control of oceanic carbon flux: the plot thickens. *Science* 280:694–696
- Azam F, Cho BC (1987) Bacterial utilization of organic matter in the sea. In: Brock T (ed) *Ecology of microbial communities*. Cambridge University Press, Cambridge, p 261–281
- Azam F, Long RA (2001) Sea snow microcosms. *Nature* 414:495–498
- Azam F, Smith DC (1991) Bacterial influence on the variability in the ocean's biogeochemical state: a mechanistic view. In: Demers S (ed) *Particle analysis in oceanography*. Springer-Verlag, New York, p 213–236
- Azam F, Martinez J, Smith DC (1993) Bacteria-organic matter coupling on marine aggregates. In: Guerrero R, Pedrós-Alí C (eds) *Trends in microbial ecology*. Spanish Society for Microbiology, Barcelona, p 410–414
- Azam F, Fonda Umani S, Funari E (1999) Significance of bacteria in the mucilage phenomenon in the northern Adriatic

- Sea. Ann Ist Super Sanità 35:411–419
- Banse K (1990) New views on the degradation and disposition of organic particles as collected by sediment traps in the ocean. Deep-Sea Res 37:1177–1195
- Bärlocher F (1992) The ecology of aquatic hyphomycetes. Ecological studies 94. Springer-Verlag, Berlin
- Becquevort S, Smith WO Jr (2001) Aggregation, sedimentation and biodegradability of phytoplankton-derived material during spring in the Ross Sea, Antarctica. Deep-Sea Res II 48:4155–4178
- Bell CR, Albright LJ (1981) Attached and free-floating bacteria in the Frazer River estuary, British Columbia, Canada. Mar Ecol Prog Ser 6:317–327
- Bell WH, Mitchell R (1972) Chemotactic and growth responses of marine bacteria to algal extracellular products. Biol Bull 143:265–277
- Bent EJ, Goulder R (1981) Planktonic bacteria in the Humber estuary: seasonal variation in population density and heterotrophic activity. Mar Biol 62:35–45
- Berger B, Hoch B, Kavka G, Herndl GJ (1996) Bacterial colonization of suspended particles in the River Danube. Aquat Microb Ecol 10:37–44
- Berman T, Viner-Mozzini Y (2001) Abundance and characteristics of polysaccharide and proteinaceous particles in Lake Kinneret. Aquat Microb Ecol 24:255–264
- Berman T, Parparov A, Simon M (1999) Carbon fluxes in limnetic seston: relative significance of respiration, solubilization and non-photosynthetic formation. Arch Hydrobiol Spec Ser Adv Limnol 54:77–90
- Bianchi M, Marty D, Teyssie JL, Fowler SW (1992) Strictly aerobic and anaerobic bacteria associated with sinking particulate matter and zooplankton fecal pellets. Mar Ecol Prog Ser 88:55–60
- Biddanda BA (1985) Microbial synthesis of macroparticulate matter. Mar Ecol Prog Ser 20:241–251
- Biddanda BA (1988) Microbial aggregation and degradation of phytoplankton-derived detritus in seawater. II. Microbial metabolism. Mar Ecol Prog Ser 42:89–95
- Biddanda BA, Pomeroy LR (1988) Microbial aggregation and degradation of phytoplankton-derived detritus in seawater. I. Microbial succession. Mar Ecol Prog Ser 42:79–89
- Bidle KD, Azam F (1999) Accelerated dissolution of diatom silica by marine bacterial assemblages. Nature 397:508–512
- Bidle KD, Azam F (2001) Bacterial control of silicon regeneration from diatom detritus: significance of bacterial ectohydrolases and species identity. Limnol Oceanogr 46:1606–1623
- Bidle KD, Fletcher M (1995) Comparison of free-living and particle-associated bacterial communities in the Chesapeake Bay by stable low-molecular weight RNA analysis. Appl Environ Microbiol 61:944–952
- Blackburn N, Azam F, Hagström Å (1997) Spatially explicit simulations of a microbial food web. Limnol Oceanogr 42:613–622
- Blackburn N, Fenchel T, Mitchell J (1998) Microscale nutrient patches in planktonic habitats shown by chemotactic bacteria. Science 282:2254–2256
- Bochdansky AB, Herndl GJ (1992) Ecology of amorphous aggregations (marine snow) in the northern Adriatic Sea. III. Zooplankton interactions with marine snow. Mar Ecol Prog Ser 87:135–146
- Böckelmann U, Manz W, Neu TR, Szewzyk U (2000) Characterization of the microbial community of lotic organic aggregates ('river snow') in the Elbe river of Germany by cultivation and molecular methods. FEMS Microbiol Ecol 33:157–170
- Brachvogel T, Schweitzer B, Simon M (2001) Dynamics and bacterial colonization of microaggregates in a large mesotrophic lake. Aquat Microb Ecol 26:23–35
- Brussaard CPD, Riegman R (1998) Influence of bacteria on phytoplankton cell mortality with phosphorus or nitrogen as the algal growth-limiting nutrient. Aquat Microb Ecol 14:271–280
- Brzezinski MA, Alldredge AL, O'Brian LM (1997) Silica cycling within marine snow. Limnol Oceanogr 42:1706–1713
- Burchard RP, Sorongon ML (1998) A gliding bacterium strain inhibits adhesion and motility of another gliding bacterium strain in a marine biofilm. Appl Environ Microbiol 64:4079–4083
- Caldwell DE, Caldwell SJ (1978) *Zoogloea ramigera* sp. associated with blooms of *Anabaena flos-aquae*. Can J Microbiol 24:922–931
- Caldwell DE, Koerber DR, Lawrence JR (1992) Confocal laser microscopy and computer image analysis. Adv Microb Ecol 12:1–67
- Capone DG, Zehr JP, Paerl HW, Bergman B, Carpenter EJ (1997) *Trichodesmium*, a globally significant marine cyanobacterium. Science 276:1221–1229
- Caron DA (1991) Heterotrophic flagellates associated with sedimenting detritus. In: Patterson DJ, Larsen J (eds) The biology of free-living heterotrophic flagellates, Spec Vol 45. Systematics Association Clarendon Press, Oxford, p 77–92
- Caron DA, Davis PG, Madin LP, Sieburth JMcN (1982) Heterotrophic bacteria and bacterivorous protozoans in oceanic macroaggregates. Science 218:795–797
- Caron DA, Davis PG, Madin LP, Sieburth JMcN (1986) Enrichment of microbial populations in macroaggregates (marine snow) from surface waters of the North Atlantic. J Mar Res 44:643–656
- Chayen J, Bitensky L, Butcher RG (1973) Practical histochemistry. John Wiley & Sons, New York
- Chen S, Eisma D (1995) Fractal geometry of *in situ* flocs in estuarine and coastal environments. Neth J Sea Res 32:173–183
- Chen S, Eisma D, Kalf J (1994) In situ size distribution of suspended matter during the tidal cycle in the Elbe estuary. Neth J Sea Res 32:37–48
- Chin WC, Orellana M, Verdugo P (1998) Spontaneous assembly of marine dissolved organic matter into polymer gels. Nature 391:568–572
- Cho BC, Azam F (1988) Major role of bacteria in biogeochemical fluxes in the ocean's interior. Nature 332:441–443
- Cole JJ (1982) Interactions between bacteria and algae in aquatic ecosystems. Annu Rev Ecol Syst 13:291–314
- Cole JJ, Likens GE, Hobbie JE (1984) Decomposition of planktonic algae in an oligotrophic lake. Oikos 42:1–9
- Costello DK, Carder KL, Betzer PR, Young RW (1989) In situ holographic imaging of settling particles: applications for individual particle dynamics and ocean flux measurements. Deep-Sea Res 36:1595–1605
- Costerton JW, Lewandowski Z, Caldwell DE, Koerber DR, Lappin-Scott HM (1995) Microbial biofilms. Annu Rev Microbiol 49:711–745
- Cottrell M, Kirchman DL (2000) Natural assemblages of marine proteobacteria and members of the *Cytophaga/Flavobacter* cluster consuming low- and high-molecular-weight dissolved organic matter. Appl Environ Microbiol 66:1692–1697
- Cowie GL, Hedges JI (1992) Sources and reactivities of amino acids in a coastal marine environment. Limnol Oceanogr 37:703–724
- Crump BC, Baross JA (1996) Particle-attached bacteria and

- heterotrophic plankton associated with the Columbia River estuarine turbidity maxima. *Mar Ecol Prog Ser* 138: 265–272
- Crump BC, Baross JA (2000) Characterization of the bacterially-active particle fraction in the Columbia River estuary. *Mar Ecol Prog Ser* 206:13–22
- Crump BC, Baross JA, Simenstad CA (1998) Dominance of particle-attached bacteria in the Columbia River estuary, USA. *Aquat Microb Ecol* 14:7–18
- Crump BC, Armbrust EV, Baross JA (1999) Phylogenetic analysis of particle-attached and free-living bacterial communities in the Columbia River, its estuary and the adjacent coastal ocean. *Appl Environ Microbiol* 65:3192–3204
- Dam HG, Drapeau DT (1995) Coagulation efficiency, organic matter glue and the dynamics of particles during a phytoplankton bloom in a mesocosm study. *Deep-Sea Res II* 42: 111–123
- Davoll PJ, Silver MW (1986) Marine snow aggregates: life history sequence and microbial community structure of abandoned larvacean houses from Monterey Bay, California. *Mar Ecol Prog Ser* 33:111–120
- Decho AW (1990) Microbial exopolymer secretion in ocean environments: their role(s) in food webs and marine processes. *Oceanogr Mar Biol Annu Rev* 28:73–153
- Degobbi D, Malej A, Fonda Umani S (1999) The mucilage phenomenon in the northern Adriatic Sea. A critical review of the present scientific hypotheses. *Ann Ist Super Sanità* 35:373–382
- Del Giorgio PA, Cole JJ (1998) Bacterial growth efficiency in natural aquatic ecosystems. *Annu Rev Ecol Syst* 29: 503–541
- DeLong EF, Franks DG, Alldredge AL (1993) Phylogenetic diversity of aggregate-attached vs. free-living marine bacterial assemblages. *Limnol Oceanogr* 38:924–934
- Dilling L, Alldredge AL (2000) Fragmentation of marine snow by swimming macrozooplankton: a new process impacting carbon cycling in the sea. *Deep-Sea Res I* 47: 1227–1245
- Dilling L, Wilson J, Steinberg D, Alldredge AL (1998) Feeding by the euphausiid *Euphausia pacifica* and the copepod *Calanus pacificus* on marine snow. *Mar Ecol Prog Ser* 170: 189–201
- Drapeau DT, Dam HG (1995) Coagulation efficiency, organic matter glues and the dynamics of particles during a phytoplankton bloom in a mesocosm study. *Deep Sea Res II* 42: 111–124
- Drapeau DT, Dam HG, Grenier G (1994) An improved flocculator design for use in particle aggregation experiments. *Limnol Oceanogr* 39:723–729
- Dugan PR, Stoner DL, Pickrum HM (1992) The genus *Zoogloea*. In: Balows A, Trüper HG, Dworkin M, Harder W, Schleifer KH (eds) *The prokaryotes*. Springer-Verlag, New York, p 3952–3964
- Eisma D (1993) Suspended matter in the aquatic environment. Springer-Verlag, New York
- Eisma D, Li A (1993) Changes in suspended matter floc size during the tidal cycle in the Dollard estuary. *Neth J Sea Res* 31:107–117
- Engel A (2000) The role of transparent exopolymer particles (TEP) in the increase in apparent particle stickiness ( $\alpha$ ) during the decline of a diatom bloom. *J Plankton Res* 22: 485–497
- Engel A, Schartau M (1999) Influence of transparent exopolymer particles (TEP) on sinking velocity of *Nitzschia closterium* aggregates. *Mar Ecol Prog Ser* 182:69–76
- Eppley RW, Harrison WG, Chisholm SW, Stewart E (1977) Particulate organic matter in surface waters off Southern California and its relation to phytoplankton. *J Mar Res* 35: 671–696
- Fell JW, Newell SY (1998) Biochemical and molecular methods for the study of marine fungi. In: Cooksey K (ed) *Molecular approaches to the study of the ocean*. Chapman & Hall, London, p 259–283
- Fenchel T (2001) Eppure si muove: many water column bacteria are motile. *Aquat Microb Ecol* 24:197–201
- Förstner U (1996) Solutes/solids interactions of metals in estuaries. *Arch Hydrobiol Spec Iss Adv Limnol* 47:271–287
- Fowler SW, GA Knauer (1986) Role of large particles in the transport of elements and organic compounds through the oceanic water column. *Prog Oceanogr* 16:147–194
- Fuhrman JA, Azam F (1980) Bacterioplankton secondary production estimates for coastal waters of British Columbia, Antarctica and California. *Appl Environ Microbiol* 39: 1085–1095
- Gerritsen J, Bradley SW (1987) Electrophoretic mobility of natural particles and cultured organisms in freshwaters. *Limnol Oceanogr* 32:1049–1058
- Gibbs RJ (1983) Effect of natural organic coatings on the coagulation of particles. *Environ Sci Technol* 17:237–240
- Glöckner FO, Fuchs BM, Amann R (1999) Bacterioplankton composition of lakes and oceans: a first comparison based on fluorescence in situ hybridization. *Appl Environ Microbiol* 65:3721–3726
- Golterman HL (1972) The role phytoplankton in detritus formation. *Mem Ist Ital Idrobiol* 29:89–103
- Gotschalk C, Alldredge AL (1989) Enhanced primary production and nutrient regeneration within aggregated marine diatoms. *Mar Biol* 103:119–129
- Grabemann I, Kappenberg J, Krause G (1996) Comparison of the dynamics of the turbidity maxima in two coastal plain estuaries. *Arch Hydrobiol Spec Iss Adv Limnol* 47: 195–205
- Graham WM, MacIntire S, Alldredge AL (2000) Diel variations of marine snow concentration in surface waters and implications for particle flux in the sea. *Deep-Sea Res I* 47: 367–395
- Green EP, Dagg MJ (1997) Mesozooplankton association with medium to large marine snow aggregates in the northern Gulf of Mexico. *J Plankton Res* 19:435–447
- Gries T, Güde H (1999) Estimates of the non-consumptive mortality of mesozooplankton by measurement of sedimentation losses. *Limnol Oceanogr* 44:459–465
- Grossart HP (1999) Interactions between marine bacteria and axenic diatoms (*Cylindrotheca fusiformis*, *Nitzschia laevis*, and *Thalassiosira weissflogii*) incubated under various conditions in the lab. *Aquat Microb Ecol* 19:1–11
- Grossart HP, Ploug H (2000) Bacterial production and growth efficiencies: direct measurements on riverine aggregates. *Limnol Oceanogr* 45:436–445
- Grossart HP, Ploug H (2001) Microbial degradation of organic carbon and nitrogen on diatom aggregates. *Limnol Oceanogr* 46:267–277
- Grossart HP, Simon M (1993) Limnetic macroscopic organic aggregates (lake snow): abundance, characteristics, and bacterial dynamics in Lake Constance. *Limnol Oceanogr* 38:532–546
- Grossart HP, Simon M (1998a) Significance of limnetic organic aggregates (lake snow) for the sinking flux of particulate organic matter in a large lake. *Aquat Microb Ecol* 15:115–125
- Grossart HP, Simon M (1998b) Bacterial colonization and microbial decomposition of limnetic organic aggregates (lake snow). *Aquat Microb Ecol* 15:127–140
- Grossart HP, Simon M, Logan BE (1997) Formation of macro-



- scopic organic aggregates (lake snow) in a large lake: the significance of transparent exopolymer particles (TEP), phyto- and zooplankton. *Limnol Oceanogr* 42:1651–1659
- Grossart HP, Berman T, Simon M, Pohlmann K (1998) Occurrence and microbial dynamics of macroscopic organic aggregates (lake snow) in Lake Kinneret, Israel, in fall. *Aquat Microb Ecol* 14:59–67
- Grossart HP, Riemann L, Azam F (2001) Bacterial motility in the sea and its ecological implications. *Aquat Microb Ecol* 25:247–258
- Guerrini F, Mazotti A, Boni L, Pistocchi A (1998) Bacterial-algal interactions in polysaccharide production. *Aquat Microb Ecol* 15:247–253
- Haake B, Ittekkot V, Ramaswamy K, Nair RR, Honjo S (1992) Fluxes of amino acids and hexosamines to the deep Arabian Sea. *Mar Chem* 40:291–314
- Haines KC, Guillard RRL (1974) Growth of vitamin B12 requiring marine diatoms with vitamin B12 producing marine bacteria. *J Phycol* 10:245–255
- Hansen JLS, Kiørboe T (1997) Quantifying interspecific coagulation efficiency of phytoplankton. *Mar Ecol Prog Ser* 159:75–79
- Hansen JLS, Timm U, Kiørboe T (1995) Adaptive significance of phytoplankton stickiness with emphasis on the diatom *Skeletonema costatum*. *Mar Biol* 123:667–676
- Hansen JLS, Kiørboe T, Alldredge AL (1996) Marine snow derived from abandoned larvacean houses: sinking rates, particle content and mechanisms of aggregate formation. *Mar Ecol Prog Ser* 141:205–215
- Heissenberger A, Leppard GG, Herndl GJ (1996) Ultrastructure of marine snow. II. Microbiological considerations. *Mar Ecol Prog Ser* 135:299–308
- Hennes KP, Suttle CA (1995) Direct counts of viruses in natural waters and laboratory cultures by epifluorescence microscopy. *Limnol Oceanogr* 40:1050–1055
- Herndl GJ (1988) Ecology of amorphous aggregations (marine snow) in the northern Adriatic Sea: II. Microbial density and activity in marine snow and its implication to overall pelagic processes. *Mar Ecol Prog Ser* 48:265–275
- Herndl GJ (1992) Marine snow in the northern Adriatic Sea: possible causes and consequences for a shallow ecosystem. *Mar Microb Food Webs* 6:149–172
- Herndl GJ, Arrieta JM, Stoderegger K (1999) Interaction between specific hydrological and microbial activity leading to extensive mucilage formation in the northern Adriatic Sea. *Ann Ist Super Sanità* 35:405–409
- Hill PS (1992) Reconciling aggregation theory with observed vertical fluxes following phytoplankton blooms. *J Geophys Res* 97:2295–2308
- Hollibaugh T, Wong PS, Murrell MC (2000) Similarity of particle-associated and free-living bacterial communities in northern San Francisco Bay, California. *Aquat Microb Ecol* 21:102–114
- Honjo S, Doherty KW, Agravall YC, Asper V (1984) Direct optical assessment of large amorphous aggregates (marine snow) in the deep ocean. *Deep-Sea Res* 31:67–76
- Hoppe HG, Ducklow HW, Karrasch B (1993) Evidence for dependency of bacterial growth on enzymatic hydrolysis of particulate organic matter in the mesopelagic ocean. *Mar Ecol Prog Ser* 93:277–283
- Huber I (1997) Bacterial diversity on lake snow: phylogenetic and in situ analyses. PhD thesis, Technische Universität München (in German)
- Imai J, Ishida Y, Nata Y (1993) Killing of marine phytoplankton by a gliding bacterium *Cytophaga* sp., isolated from the coastal sea of Japan. *Mar Biol* 116:527–532
- Jackson GA (1990) A model of the formation of the marine algal flocs by physical coagulation processes. *Deep-Sea Res* 37:1197–1211
- Jackson GA (1994) Particle trajectories in rotating cylinders: implications for aggregation incubations. *Deep-Sea Res I* 41:429–437
- Jackson GA (1995) Comparing observed changes in particle size spectra with those predicted using coagulation theory. *Deep-Sea Res* 42:159–184
- Jackson GA (2001) Effect of coagulation on a model planktonic food web. *Deep Sea Res I* 48:95–123
- Jackson GA, Maffione R, Costello DK, Alldredge AL, Logan BE, Dam HG (1997) Particle size spectra between 1 µm and 1 cm at Monterey Bay determined using multiple instruments. *Deep-Sea Res I* 44:1739–1767
- Jähmlich S, Thomsen L, Graf G (1999) Factors controlling aggregate formation in the benthic boundary layer of the Mecklenburg Bight (western Baltic Sea). *J Sea Res* 41:245–254
- Johannes RE (1968) Nutrient regeneration in lakes and ocean. *Adv Microbiol Sea* 1:203–213
- Johnson BD, Wangersky PJ (1985) A recording backscattering meter and camera system for examination of the distribution and morphology of macroaggregates. *Deep-Sea Res* 32:1143–1150
- Johnson BD, Zhou X, Wangersky PJ (1986) Surface coagulation in seawater. *Neth J Sea Res* 20:201–210
- Johnson BD, Kranck K, Muschenheim DK (1994) Physicochemical factors in particle aggregation. In: Wotton RS (ed) *The biology of particles in aquatic systems*. Lewis Publishers, Boca Raton, FL, p 75–96
- Jürgens K, Güde H (1994) The potential importance of grazing-resistant bacteria in planktonic systems. *Mar Ecol Prog Ser* 112:169–188
- Kaltenböck E, Herndl GJ (1992) Ecology of amorphous aggregations (marine snow) in the northern Adriatic Sea. IV. Dissolved nutrients and the autotrophic community associated with marine snow. *Mar Ecol Prog Ser* 87:147–159
- Kamjunke N, Mehner T (2001) Coupling the microbial food web with fish: are bacteria attached to cyanobacteria an important food source for underyearling roach? *Freshw Biol* 46:633–639
- Kamjunke N, Schmidt K, Pflugmacher S, Mehner T (2002) Consumption of cyanobacteria by roach (*Rutilus rutilus*): useful or harmful to the fish? *Freshw Biol* 47:243–254
- Karl DM, Tilbrook BD (1994) Production and transport of methane in oceanic particulate organic matter. *Nature* 368:732–734
- Karl DM, Knauer GA, Martin JH (1988) Downward flux of particulate organic matter in the ocean: a particle decomposition paradox. *Nature* 332:438–441
- Karner M, Herndl GJ (1992) Extracellular enzymatic activity and secondary production in free-living and marine snow associated bacteria. *Mar Biol* 113:341–347
- Kawano Y, Nagawa Y, Nakanishi H, Nakajima H, Matsuo M, Higashihara T (1997) Production of thiotropocin by a marine bacterium, *Caulobacter* sp. and its antimicrobial activities. *J Mar Biotechnol* 5:225–229
- Kepkay PE (1994) Particle aggregation and the biological reactivity of colloids. *Mar Ecol Prog Ser* 109:293–304
- Kerner M, Gramm H (1995) Changes in oxygen consumption at the sediment-water interface formed by settling seston from the Elbe estuary. *Limnol Oceanogr* 40:544–555
- Kerner M, Krogmann D (1994) Partitioning of trace metals in suspended matter from the Elbe estuary fractionated by a sedimentation method. *Neth J Sea Res* 33:19–27
- Kilps RJ, Logan BE, Alldredge AL (1994) Fractal dimensions of marine snow determined from image analysis of in situ

- photographs. *Deep-Sea Res I* 41:1159–1169
- Kjørboe T (2000) Colonization of marine snow aggregates by invertebrate zooplankton: abundance, scaling and possible role. *Limnol Oceanogr* 45:479–484
- Kjørboe T (2001) Formation and fate of marine snow: small-scale processes with large-scale implications. *Sci Mar* 65(Suppl 2):57–71
- Kjørboe T, Hansen JLS (1993) Phytoplankton aggregate formation: observations of patterns and mechanisms of cell sticking and the significance of exopolymer material. *J Plankton Res* 15:993–1018
- Kjørboe T, Jackson GA (2001) Marine snow, organic solute plumes, and optimal chemosensory behavior of bacteria. *Limnol Oceanogr* 46:1309–1318
- Kjørboe T, Thygesen UH (2001) Fluid motion and solute distribution around sinking aggregates. II. Implications for remote detection by colonizing zooplankters. *Mar Ecol Prog Ser* 211:15–25
- Kjørboe T, Andersen KP, Dam HG (1990) Coagulation efficiency and aggregate formation in marine phytoplankton. *Mar Ecol Prog Ser* 107:235–245
- Kjørboe T, Lundsgaard C, Olesen M, Hansen JLS (1994) Aggregation and sedimentation processes during a spring phytoplankton bloom: a field experiment to test coagulation theory. *J Mar Res* 52:297–323
- Kjørboe T, Hansen, JLS, Alldredge AL, Jackson GA and 5 others (1996) Sedimentation of phytoplankton during a diatom bloom: rates and mechanisms. *J Mar Res* 54: 1123–1148
- Kjørboe T, Tiselius P, Mitchell-Innes B, Hansen JLS, Visser AW, Mari X (1998) Intensive aggregate formation with low vertical flux during an upwelling-induced diatom bloom. *Limnol Oceanogr* 43:104–116
- Kjørboe T, Ploug H, Thygesen UH (2001) Fluid motion and solute distribution around sinking aggregates. I. Small scale fluxes and heterogeneity of nutrients in the pelagic environment. *Mar Ecol Prog Ser* 211:1–13
- Kirchman DL (1993) Particulate detritus and bacteria in marine environments. In: Ford TE (ed) *Aquatic microbial ecology*. Blackwell Scientific, Oxford, p 321–342
- Kirchman DL, Ducklow HW (1987) Trophic dynamics of particle-bound bacteria in aquatic ecosystems: a review. In: Moriarty DJW, Pullin RSV (eds) *Detritus and microbial ecology in aquaculture*. ICLARM conference proceedings, 14:54–82
- Kirchman DL, K'nees E, Hodson RE (1985) Leucine incorporation and its potential as a measure of protein synthesis by bacteria in natural systems. *Appl Environ Microbiol* 49: 599–607
- Knoll S, Zwisler W, Simon M (2001) Bacterial colonization of early stages of limnetic diatom microaggregates. *Appl Environ Microbiol* 25:141–150
- Kochert G, Olson LW (1970) Endosymbiotic bacteria in *Volvox carteri*. *Trans Am Microsc Soc* 89:475–478
- Kodama M, Ogata T, Sato T, Sakamoto S (1990) Possible association of marine bacteria with paralytic shellfish toxicity of bivalves. *Mar Ecol Prog Ser* 61:203–206
- Koike I, Shigemitsu H, Kazuki T, Kogure K (1990) Role of sub-micrometer particles in the ocean. *Nature* 345:242–244
- Lampitt RS (1996) Snow falls in the ocean. In: Summerhayes CP, Thorpe SA (eds) *Oceanography an illustrated guide*. Manson Publishing, London, 96–112
- Lampitt RS, Noji T, v Bodungen B (1990) What happens to zooplankton fecal pellets? *Mar Biol* 104:15–23
- Lampitt RS, Wishner KF, Turley CM, Angel MV (1993a) Marine snow studies in the North Atlantic Ocean: distribution, composition and role as a food source for migrating zooplankton. *Mar Biol* 116:689–702
- Lampitt RS, Hillier WR, Challenor PG (1993b) Seasonal and diel variation in the open ocean concentration of marine snow aggregates. *Nature* 362:737–739
- Larson ET, Shanks AL (1996) Consumption of marine snow by two species of juvenile mullet and its contribution to their growth. *Mar Ecol Prog Ser* 130:19–28
- Lee N, Nielsen PK, Andreassen KH, Juretschko S, Nielsen JL, Schleifer KH, Wagner M (1999) Combination of fluorescent in situ hybridization and microautoradiography—a new tool for structure-function analysis in microbial ecology. *Appl Environ Microbiol* 65:1289–1297
- Leppard CG, Massalski A, Lean DRS (1979) Electron-opaque microscopic fibrils in lakes: their demonstration, their biological derivation and their potential significance in the redistribution of cations. *Protoplasma* 92:289–309
- Li X, Logan BE (1995) Size distribution and fractal geometry of particles during a simulated diatom bloom in a mesocosm. *Deep-Sea Res II* 42:125–138
- Lochte K, Turley CM (1988) Bacteria and cyanobacteria associated with phytodetritus in the deep sea. *Nature* 333: 67–69
- Logan BE, Alldredge AL (1989) Potential for increased nutrient uptake by flocculating diatoms. *Mar Biol* 101:443–450
- Logan BE, Hunt JR (1987) Advantages to microbes of growth in permeable aggregates in marine systems. *Limnol Oceanogr* 32:1034–1048
- Logan BE, Kilps JR (1995) Fractal dimensions of aggregates formed in different fluid mechanical environments. *Water Res* 29:443–453
- Logan BE, Wilkinson DB (1990) Fractal geometry of marine snow and other biological aggregates. *Limnol Oceanogr* 35:130–136
- Logan BE, Grossart HP, Simon M (1994) Direct observation of phytoplankton, TEP and aggregates on polycarbonate filters using brightfield microscopy. *J Plankton Res* 16: 1811–1815
- Logan BE, Passow U, Alldredge AL, Grossart HP, Simon M (1995) Mass sedimentation of diatom blooms as large aggregates is driven by coagulation of transparent exopolymer particles (TEP). *Deep-Sea Res II* 42:203–214
- Long RA, Azam F (1996) Abundant protein-containing particles in the sea. *Aquat Microb Ecol* 10:213–221
- Long RA, Azam F (2001) Antagonistic interactions among marine pelagic bacteria. *Appl Environ Microbiol* 67: 4975–4983
- Manz W, Wendt-Potthoff K, Neu TR, Szewzyk U (1999) Phylogenetic composition, spatial structure, and dynamics of lotic biofilms investigated by fluorescent in situ hybridization and confocal laser scanning microscopy. *Microb Ecol* 37:225–237
- Mari X (1999) Carbon content and C:N ratio of transparent exopolymer particles (TEP) produced by bubbling exudates of diatoms. *Mar Ecol Prog Ser* 183:59–71
- Mari X, Burd A (1998) Seasonal size spectra of transparent exopolymer particles (TEP) in a coastal sea and comparison with those predicted using coagulation theory. *Mar Ecol Prog Ser* 163:63–76
- Mari X, Kjørboe T (1996) Abundance, size distribution and bacterial colonization of transparent exopolymer particles (TEP) during spring in the Kattegatt. *J Plankton Res* 18: 969–986
- Mari X, Beauvais S, Lemée R, Pedrotti ML (2001) Non-Redfield C:N ratio of transparent exopolymer particles in the northwestern Mediterranean Sea. *Limnol Oceanogr* 46: 1831–1836
- Marty DG (1993) Methanogenic bacteria in seawater. *Limnol*

- Oceanogr 38:452–456
- McCave IN (1984) Size spectra and aggregation of suspended particles in the deep ocean. *Deep-Sea Res* 31:329–352
- McIntyre S, Alldredge AL, Gotschalk C (1995) Accumulation of marine snow at density discontinuities in the water column. *Limnol Oceanogr* 40:449–468
- Melchiorri-Santolini U, Hopton JW (1972) Detritus and its role in aquatic ecosystems. *Mem Ist Ital Idrobiol (Spec)* 29
- Milligan TG (1995) An examination of the settling behavior of a flocculated suspension. *Neth J Sea Res* 32:163–171
- Mitchell JG, Pearson L, Dillon S, Kantalis K (1995) Natural assemblages of marine bacteria exhibiting high-speed motility and large accelerations. *Appl Environ Microbiol* 62:4436–4440
- Moeseneder M, Winter C, Herndl GJ (2001) Horizontal and vertical complexity of attached and free-living bacteria in the eastern Mediterranean Sea, determined by 16S rDNA and 16S rRNA fingerprints. *Limnol Oceanogr* 46:95–107
- Mopper K, Zhou J, Ramana KS, Passow U, Dam H, Drapeau D (1995) The role of surface-active carbohydrates in the flocculation of a diatom bloom in a mesocosm. *Deep-Sea Res II* 42:47–73
- Mostajir B, Dolan JR, Rassoulzadegan R (1995) Seasonal variations of pico- and nano-detrital particles (DAPI yellow particles, DYP) in the Ligurian Sea (NW Mediterranean). *Aquat Microb Ecol* 9:267–277
- Müller-Niklas G, Schuster G, Kaltenböck E, Herndl GJ (1994) Organic content and bacterial metabolism in amorphous aggregations in the northern Adriatic Sea. *Limnol Oceanogr* 39:58–68
- Neu TR (2000) *In situ* cell and glycoconjugate distribution in river snow studied by confocal laser scanning microscopy. *Aquat Microb Ecol* 21:85–95
- O'Melia CR (1987) Particle-particle interactions. In: Stumm W (ed) *Aquatic surface chemistry*. Wiley-Interscience, New York, p 385–403
- Oremland RS (1979) Methanogenic activity in plankton samples and fish intestines — mechanisms for in situ methanogenesis in oceanic surface waters. *Limnol Oceanogr* 24:1136–1141
- Ouverney CC, Fuhrman JA (1999) Combined microautoradiography-16S rRNA probe technique for determination of radioisotope uptake by specific microbial cell types in situ. *Appl Environ Microbiol* 65:1746–1752
- Paerl HW (1974) Bacterial uptake of dissolved organic matter in relation to detrital aggregation in marine and freshwater systems. *Limnol Oceanogr* 17:966–972
- Paerl HW, Pinckney JL (1996) A minireview of microbial consortia: their roles in aquatic production and biogeochemical cycling. *Microb Ecol* 31:225–247
- Parparov A, Berman T, Grossart HP, Simon M (1998) The metabolic activity of seston-associated microorganisms in Lake Kinneret. *Aquat Microb Ecol* 15:77–78
- Passow U (2000) Formation of transparent exopolymer particles, TEP, from dissolved precursor material. *Mar Ecol Prog Ser* 192:1–11
- Passow U, Alldredge AL (1994) Distribution, size and bacterial colonization of transparent exopolymer particles (TEP) in the ocean. *Mar Ecol Prog Ser* 113:185–198
- Passow U, Alldredge AL (1995) Mass aggregation of a diatom bloom in a mesocosm. *Deep-Sea Res II* 42:99–110
- Passow U, Alldredge AL (2000) Do transparent exopolymer particles (TEP) inhibit grazing by the euphausiid *Euphausia pacifica*? *J Plankton Res* 21:2203–2217
- Passow U, Alldredge AL, Logan BE (1994) The role of particulate carbohydrate exudates in the flocculation of diatom blooms. *Deep Sea Res I* 41:335–357
- Phillips CJ, Smith Z, Embley TM, Prosser JI (1999) Phylogenetic differences between particle-associated and planktonic ammonia-oxidizing bacteria of the  $\beta$ -subdivision of the class Proteobacteria in the northwestern Mediterranean Sea. *Appl Environ Microbiol* 65:779–786
- Pilskaln CH, Lehmann C, Paduan JB, Silver MW (1998) Spatial and temporal dynamics in marine aggregate abundance, sinking rate and flux: Monterey Bay, central California. *Deep-Sea Res II* 45:1803–1837
- Ploug H (2001) Small-scale oxygen fluxes and remineralization in sinking aggregates. *Limnol Oceanogr* 46:1624–1631
- Ploug H, Grossart HP (1999) Bacterial production and respiration in suspended aggregates — a matter of the incubation method. *Aquat Microb Ecol* 20:21–29
- Ploug H, Grossart HP (2000) Bacterial growth and grazing on diatom aggregates: respiratory carbon turnover as a function of aggregate size and sinking velocity. *Limnol Oceanogr* 45:1467–1475
- Ploug H, Jørgensen BB (1999) A net-jet flow system for mass transfer and microelectrode studies in sinking aggregates. *Mar Ecol Prog Ser* 176:270–290
- Ploug H, Kühl M, Buchholz-Cleven B, Jørgensen BB (1997) Anoxic aggregates — an ephemeral phenomenon in the pelagic environment. *Aquat Microb Ecol* 13:285–294
- Ploug H, Grossart HP, Azam F, Jørgensen BB (1999) Photosynthesis, respiration and carbon turnover in sinking marine snow from surface waters of Southern California Bight: implications for the carbon cycling in the ocean. *Mar Ecol Prog Ser* 179:1–11
- Pomeroy LR, Deibel D (1980) Aggregation of organic matter by pelagic tunicates. *Limnol Oceanogr* 25:643–652
- Pomeroy LR, Hanson RB, McGillivray PA, Sherr BF, Kirchman D, Deibel D (1984) Microbiology and chemistry of fecal products of pelagic tunicates: rates and fates. *Bull Mar Sci* 35:426–439
- Pringsheim EG (1912) Die Kultur von Algen in Agar. *Beitr Biol Pflanz* 11:305–333
- Proctor LM, Fuhrman JA (1991) Roles of virus infection in organic particle flux. *Mar Ecol Prog Ser* 69:133–142
- Rath J, Wu KY, Herndl GJ, DeLong EF (1998) High phylogenetic diversity in a marine snow-associated bacterial assemblage. *Aquat Microb Ecol* 14:261–269
- Rausch de Traubenberg, C, Lassus P (1991) Dinoflagellate toxicity: are marine bacteria involved? Evidence from the literature. *Mar Microb Food Webs* 5:205–226
- Reichenbach H (1992) The order *Cytophagales*. In: Balows A, Trüper HG, Dworkin M, Harder W, Schleifer KH (eds) *The prokaryotes*. Springer-Verlag, New York, p 3631–3675
- Riebesell U (1991a) Particle aggregation during a diatom bloom. I. Physical aspects. *Mar Ecol Prog Ser* 69:273–280
- Riebesell U (1991b) Particle aggregation during a diatom bloom. II. Biological aspects. *Mar Ecol Prog Ser* 69:281–291
- Riebesell U (1992) The formation of large marine snow and its sustained residence in surface waters. *Limnol Oceanogr* 37:63–76
- Riebesell U, Schloss I, Smetacek V (1991) Aggregation of algae released from melted sea ice: implication for seeding and sedimentation. *Polar Biol* 11:239–248
- Rogerson A, Laybourn-Parry J (1992) Aggregate dwelling protozooplankton communities in estuaries. *Arch Hydrobiol* 125:411–422
- Schäfer B, Harms H (1996) Occurrence of different ammonia-oxidizing bacteria in the River Elbe between Hamburg and Glückstadt. *Arch Hydrobiol Spec Iss Adv Limnol* 47:141–148
- Schumann R, Rentsch D (1998) Staining particulate organic matter with DTAF — a fluorescence dye for carbohydrates

- and protein: a new approach and application of a 2D image analysis system. *Mar Ecol Prog Ser* 163:77–88
- Schuster S, Herndl GJ (1995) Formation and significance of transparent exopolymeric particles in the northern Adriatic Sea. *Mar Ecol Prog Ser* 124:227–236
- Schweitzer B, Huber I, Amann R, Ludwig W, Simon M (2001) Alpha- and beta-proteobacteria control the consumption and release of amino acids on lake snow aggregates. *Appl Environ Microbiol* 67:632–645
- Shanks AL, Del Carmen KA (1997) Larval polychaetes are strongly associated with marine snow. *Mar Ecol Prog Ser* 154:211–221
- Shanks AL, Edmondson EW (1989) Laboratory-made artificial marine snow: a biological model of the real thing. *Mar Biol* 101:463–470
- Shanks AL, Edmondson EW (1990) Vertical flux of metazoans (holoplankton, meiofauna, and larval invertebrates) due to their association with marine snow. *Limnol Oceanogr* 35:455–463
- Shanks AL, Reeder ML (1993) Reducing microzones and sulfide production in marine snow. *Mar Ecol Prog Ser* 96:43–47
- Shanks AL, Trent JD (1979) Marine snow: microscale nutrient patches. *Limnol Oceanogr* 24:43–47
- Shanks AL, Walters K (1996) Feeding by a heterotrophic dinoflagellate (*Noctiluca scintillans*) in marine snow. *Limnol Oceanogr* 41:177–181
- Shanks AL, Walters K (1997) Holoplankton, meroplankton, and meiofauna associated with marine snow. *Mar Ecol Prog Ser* 156:75–86
- Sieburth JMcN (1968) The influence of algal antibiosis on the ecology of marine microorganisms. In: Droop S, Wood F (eds) *Advances in microbiology of the sea*. Academic Press, London, p 217–233
- Silver MW, Shanks AL, Trent JD (1978) Marine snow: microplankton habitat and source of small scale patchiness in pelagic populations. *Science* 201:371–373
- Silver MW, Coale SL, Pilskaln CH, Steinberg DR (1998) Giant aggregates: importance as microbial centers and agents of material flux in the mesopelagic zone. *Limnol Oceanogr* 43:498–507
- Simon M (1987) Biomass and production of small and large free-living and attached bacteria in Lake Constance. *Limnol Oceanogr* 32:591–607
- Simon M, Azam F (1989) Protein content and protein synthesis rates of planktonic marine bacteria. *Mar Ecol Prog Ser* 51:201–213
- Simon M, Alldredge AL, Azam F (1990) Bacterial carbon dynamics on marine snow. *Mar Ecol Prog Ser* 65:205–211
- Simon M, Welschmeyer NA, Kirchman DL (1992) Bacterial production and the sinking flux of particulate organic matter in the subarctic Pacific. *Deep-Sea Res* 39:1997–2008
- Simon M, Glöckner FO, Amann R (1999) Different community structure and temperature optima of heterotrophic picoplankton in various regions of the Southern Ocean. *Aquat Microb Ecol* 18:275–284
- Simon M, Jontofsohn M, Parparov A, Berman T (2000) Turnover of combined amino acids and carbohydrates on organic aggregates and in the bulk water in Lake Kinneret and other pelagic ecosystems. *Arch Hydrobiol Spec Iss Adv Limnol* 55:365–377
- Smetacek V (1985) Role of sinking in diatom life history cycles: ecological, evolutionary and geological significance. *Mar Biol* 84:239–251
- Smith DC, Simon M, Alldredge AL, Azam F (1992) Intense hydrolytic enzyme activity on marine aggregates and implications for rapid particle dissolution. *Nature* 359:139–142
- Smith DC, Steward GF, Long RA, Azam F (1995) Bacterial mediation of carbon fluxes during a diatom bloom in a mesocosm. *Deep-Sea Res* 42:75–97
- Stal LJ, Staal M, Villbrandt M (1999) Nutrient control of cyanobacterial blooms in the Baltic Sea. *Aquat Microb Ecol* 18:165–173
- Steemann-Nielsen E (1955) The production of antibiotics by planktonic algae and its effect on bacterial activities in the sea. *Deep-Sea Res* 3:281–286
- Stehr G, Zörner S, Böttcher B, Koops HP (1995) Exopolymers: an ecological characteristic of a floc-attached ammonium oxidizing bacteria. *Microb Ecol* 30:115–126
- Steinberg DK, Silver MW, Pilskaln CH, Coale SL, Paduan JB (1994) Midwater zooplankton communities on pelagic detritus (giant larvacean houses) in Monterey Bay, California. *Limnol Oceanogr* 39:1606–1620
- Steinberg DK, Silver MW, Pilskaln CH (1997) Role of mesopelagic zooplankton in the community metabolism of giant larvacean house detritus in Monterey Bay, California, USA. *Mar Ecol Prog Ser* 147:167–179
- Stoderegger KE, Herndl GJ (1998) Production and release of bacterial capsular material and its subsequent utilization by marine bacterioplankton. *Limnol Oceanogr* 43:877–884
- Stoderegger KE, Herndl GJ (1999) Production of exopolymer particles by marine bacterioplankton under contrasting turbulence conditions. *Mar Ecol Prog Ser* 189:9–16
- Stolzenbach KD (1993) Scavenging of small particles by fast-sinking porous aggregates. *Deep Sea Res* 40:359–369
- Stumm W, Sigg L, Sulzberger B (1994) The role of coordination at the surface of aquatic particles. In: Buffle J, de Vitre RR (ed) *Chemical and biological regulation of aquatic ecosystems*. Lewis Publishers, Boca Raton, FL, p 43–88
- Suzuki N, Kato K (1953) Studies on suspended materials. Marine snow in the sea. I. Sources of marine snow. *Bull Fac Fish Hokkaido Univ* 4:132–135
- Thiel H, Pfannkuche O, Schriever G, Lochte K and 6 others (1988) Phytodetritus on the deep-sea floor in a central oceanic region of the Northeast Atlantic. *Biol Oceanogr* 6:203–239
- Thomsen L, Graf G (1994) Boundary layer characteristics of the continental margin of the western Greenland Sea. *Oceanol Acta* 17:597–607
- Tiselius P, Kiørboe T (1998) Colonization of diatom aggregates by the dinoflagellate *Noctiluca scintillans*. *Limnol Oceanogr* 43:154–159
- Tiselius P, Kuylensstierna M (1996) Growth and decline of a diatom spring bloom: phytoplankton species composition, formation of marine snow and the role of heterotrophic dinoflagellates. *J Plankton Res* 18:133–155
- Turley CM, Mackie PJ (1994) Biogeochemical significance of attached and free-living bacteria and the flux of particles in the NE Atlantic Ocean. *Mar Ecol Prog Ser* 115:191–203
- Turley CM, Stutt ED (2000) Depth-related cell-specific bacterial leucine incorporation rates on particles and its biogeochemical significance in the Northwest Mediterranean. *Limnol Oceanogr* 45:419–425
- Turner JT (2002) Zooplankton fecal pellets, marine snow and sinking phytoplankton blooms. *Aquat Microb Ecol* 27:57–102
- Unanue M, Azua I, Arrieta JM, Herndl GJ, Irriberri J (1998a) Laboratory-made particles as a useful approach to analyze microbial processes in marine macroaggregates. *FEMS Microbiol Ecol* 26:325–334
- Unanue M, Azua I, Arrieta JM, Labirua-Iturburu A, Egea L, Irriberri J (1998b) Bacterial colonization and ectoenzymatic activity in phytoplankton-derived model particles. Cleavage of peptides and uptake of amino acids. *Microb*



- Ecol 35:136–146
- Wagner M, Amann R, Lemmer H, Schleifer KH (1993) Probing activated sludge with oligonucleotides specific for *Proteobacteria*: inadequacy of culture-dependent methods for describing microbial community structure. Appl Environ Microbiol 59:1520–1525
- Wagner M, Amann R, Kämpfer P, Assmus B, Hartmann A, Hutzler P, Springer N, Schleifer KH (1994) Identification and in situ detection of gram-negative filamentous bacteria in activated sludge. Syst Appl Microbiol 17:405–417
- Waite A, Gallager S, Dam HG (1997) New measurements of phytoplankton aggregation in a flocculator using videography and image analysis. Mar Ecol Prog Ser 155:77–88
- Wakeham SG, Lee C (1993) Production, transport and alteration of particulate organic matter in the marine water column. In: Engel MH, Macko SA (eds) Organic geochemistry. Plenum Press, New York, p 145–169
- Walsh ID, Gardner WD (1992) Comparison of large particle camera profiles with sediment trap fluxes. Deep-Sea Res 39:1817–1834
- Walsh ID, Gardner WD, Richardson MJ, Chung SP, Plattner CA, Asper VL (1997) Particle dynamics as controlled by the flow field of the eastern equatorial Pacific. Deep-Sea Res II 44:2025–2047
- Weiss P, Schweitzer B, Amann R, Simon M (1996) Identification *in situ* and dynamics of bacteria on limnetic organic aggregates (Lake Snow). Appl Environ Microbiol 62: 1998–2005
- Wells ML, Goldberg ED (1991) Occurrence of small colloids in sea water. Nature 353:324–344
- Winstanley C, Morgan JAW (1997) The bacterial flagellin gene as a biomarker for detection, population genetics and epidemiological analysis. Microbiology 143: 3071–3084
- Wommack KE, Colwell RR (2000) Virioplankton: viruses in aquatic ecosystems. Microbiol Mol Biol Rev 64:69–114
- Worm J, Søndergaard M (1998a) Dynamics of heterotrophic bacteria attached to *Mycrocystis* spp. (Cyanobacteria). Aquat Microb Ecol 14:18–28
- Worm J, Søndergaard M (1998b) Alcian Blue-stained particles in a eutrophic lake. J Plankton Res 20:179–186
- Wotton RS (1994) The biology of particles in aquatic systems. Lewis Publishers, Boca Raton, FL
- Zhou J, Mopper K, Passow U (1998) The role of surface-active carbohydrates in the formation of transparent exopolymer particles by bubble adsorption of seawater. Limnol Oceanogr 43:1860–1871
- Zimmermann H (1997) The microbial community on aggregates in the Elbe estuary. Aquat Microb Ecol 13:37–46
- Zimmermann H, Kausch H (1996) Microaggregates in the Elbe Estuary: structure and colonization during spring. Arch Hydrobiol Spec Iss Adv Limnol 48:85–92
- Zimmermann-Timm H, Holst H, Müller S (1998) Seasonal dynamics of aggregates and their typical biocoenosis in the Elbe Estuary. Estuaries 21:613–621
- ZoBell CE (1941) Studies on marine bacteria. I. The cultural requirements of heterotrophic aerobes. J Mar Res 4:42–75
- Zutic V, Tomaic J (1988) On the formation of organic coatings on marine particles: interactions of organic matter at hydrous aluminum/seawater interfaces. Mar Chem 23: 51–67
- Zwisler W, Simon M (2002) Seasonal patterns of the bacterioplankton community composition in a large mesotrophic lake. Aquat Microb Ecol (in press)

Editorial responsibility: Karin Lochte,  
Kiel, Germany

Submitted: November 27, 2000; Accepted: April 2, 2002  
Proofs received from author(s): May 22, 2002