

Singularities in ternary mixtures of k -core percolation

Davide Cellai and James P. Gleeson

Abstract Heterogeneous k -core percolation is an extension of a percolation model which has interesting applications to the resilience of networks under random damage. In this model, the notion of node robustness is local, instead of global as in uniform k -core percolation. One of the advantages of k -core percolation models is the validity of an analytical mathematical framework for a large class of network topologies. We study ternary mixtures of node types in random networks and show the presence of a new type of critical phenomenon. This scenario may have useful applications in the stability of large scale infrastructures and the description of glass-forming systems.

1 Introduction

Percolation, with its many modifications and extensions, is a problem with a venerable past and many applications in the most diverse disciplines [18]. With the rapid development of network science, several new problems and models have been introduced. Among the different directions, k -core percolation constitutes a development which, in spite of its somewhat simple definition, is able to encapsulate a number of interesting problems which can often be approached with a robust mathematical formalism. Given a network, a k -core is defined as the subnetwork where each node has at least k neighbours in the same subnetwork. A k -core can equivalently be defined as the subnetwork remaining after a culling process consisting in recursively removing all the nodes with degree lower than k . k -core percolation has

Davide Cellai

MACSI, Department of Mathematics and Statistics, University of Limerick, Ireland, e-mail: davide.cellai@ul.ie

James P. Gleeson

MACSI, Department of Mathematics and Statistics, University of Limerick, Ireland, e-mail: james.gleeson@ul.ie

applications in many different disciplines including jamming [15], neural networks [8], granular gases [1], evolution [14], social sciences [13] and the metal-insulator transition [4]. The application of k -core percolation we are interested in here, however, refers to the stability of a network under random damage. A fraction $(1 - p)$ of nodes is removed (together with the adjacent edges) and the size M_k of the largest k -core cluster is studied as a function of p . The k -core strength M_k can vanish either continuously or discontinuously, a scenario that can be described in terms of phase transitions (second and first order, respectively) and can be important in the stability of large scale infrastructures [9, 10]. A recent extension of k -core percolation, named heterogeneous k -core (HKC) percolation considers the threshold k_i as a local property, and allows therefore the mixing of different critical phenomena. An analytical formalism has been introduced to approach this model on networks and binary mixtures of thresholds k have been investigated [2, 5, 6]. On a wide class of networks, this model displays phase diagrams which are topologically equivalent to the ones calculated for a recently introduced spin model of glass-forming systems [17, 16]. This model can be seen as a heterogeneous development of the Fredrickson-Andersen (FA) model of facilitated spins, where the facilitation consists of a local constraint in the number of spins down in order for the considered spin to be able to flip [11, 17]. It has been shown that this model reproduces characteristic signatures of glass-forming systems called *glass-transition singularities*, which correspond to distinctive critical phenomena for the appropriate choice of the parameters [12, 16]. One of the glass-transition singularities which has not yet been explored is the so-called A_4 singularity, characterised by the coalescence of two critical points. As the HKC model appears to reproduce all the relevant critical phenomena for binary mixtures, it is interesting to look for this singularity as well. The purpose of this paper, therefore, is to investigate, for the first time, ternary mixtures in HKC percolation and calculate the critical point corresponding to an A_4 singularity.

In Section 2 we present the analytical formalism used here (from [2]), in Section 3 we sketch the behaviour of the model for binary mixtures, in Section 4 we show a case of ternary mixture with an A_4 singularity and in Section 5 we give the conclusions.

2 Formalism

Let k_i be the k -core threshold of the node i . A fraction $(1 - p)$ of nodes are randomly removed: the problem consists in calculating the size of the HKC (if it exists). We consider the configuration model of random networks, defined as the maximally random network with a given degree distribution $P(q)$. The configuration model has the property of being locally tree-like, i. e. the number of finite loops vanishes for infinite networks. This property allows to consider, in an infinite network, the HKC equivalent to the $(k_i - 1)$ -ary subtree, defined as the tree in which, as we traverse it, each encountered vertex has at least $k_i - 1$ child edges. Then we can write a self-

consistent equation for Z , the probability that a randomly chosen node is the root of a $(k_i - 1)$ -ary subtree [2]:

$$\begin{aligned} Z = & pr \sum_{q=k_a}^{\infty} \frac{qP(q)}{\langle q \rangle} \sum_{l=k_a-1}^{q-1} \binom{q-1}{l} Z^l (1-Z)^{q-1-l} + \\ & + ps \sum_{q=k_b}^{\infty} \frac{qP(q)}{\langle q \rangle} \sum_{l=k_b-1}^{q-1} \binom{q-1}{l} Z^l (1-Z)^{q-1-l} + \\ & + p(1-r-s) \sum_{q=k_c}^{\infty} \frac{qP(q)}{\langle q \rangle} \sum_{l=k_c-1}^{q-1} \binom{q-1}{l} Z^l (1-Z)^{q-1-l}. \end{aligned} \quad (1)$$

where the three thresholds $\mathbf{k} = (k_a, k_b, k_c)$ are randomly assigned to nodes with probability r , s and $(1-r-s)$, respectively. In this paper we always assume $k_i \geq 2$, so we do not need to consider the case where there may be finite clusters in the HKC. Due to the absence of finite loops, no finite HKC clusters can exist if $k_i \geq 2$ for all nodes i [10, 2].

We can then write the probability M_{abc} that a randomly chosen node is in the HKC, for a mixture of three types of nodes:

$$\begin{aligned} M_{abc}(p) = & pr \sum_{q=k_a}^{\infty} P(q) \sum_{l=k_a}^q \binom{q}{l} Z^l (1-Z)^{q-l} + \\ & ps \sum_{q=k_b}^{\infty} P(q) \sum_{l=k_b}^q \binom{q}{l} Z^l (1-Z)^{q-l} + \\ & p(1-r-s) \sum_{q=k_c}^{\infty} P(q) \sum_{l=k_c}^q \binom{q}{l} Z^l (1-Z)^{q-l}. \end{aligned} \quad (2)$$

3 Binary mixtures in heterogeneous k -core percolation

In homogeneous k -core percolation it is known that, for networks with a fast decreasing degree distribution, namely $P(q) < 1/q^\gamma$ with $\gamma > 3$ for $q \rightarrow \infty$, a randomly damaged network collapses continuously for $k \leq 2$ and discontinuously for $k \geq 3$ [7, 3, 10]. In recent papers, it has also been shown that in the case of binary mixtures HKC percolation is characterized by a few different critical phenomena. For example, in mixtures where the two values of k are associated to continuous and discontinuous transitions, respectively, a critical point (also called A_3 singularity or cusp singularity) is usually observed [2, 6]. That is the case, for instance, of the mixtures $\mathbf{k} = (1, 3)$ or $\mathbf{k} = (2, 4)$, where the line of first order transitions ends in a critical point and a line of second order transition intersects the former. The case $\mathbf{k} = (2, 3)$, though, is quite peculiar as the critical line exactly matches the line of first order transitions giving rise to a tricritical point (Fig. 1) [5].




Fig. 1 Comparison of the phase diagrams of the cases $\mathbf{k} = (1, 3)$, $\mathbf{k} = (2, 3)$ and $\mathbf{k} = (2, 4)$ for Erdős-Rényi graphs with $z_1 = 10$. Continuous lines represent second order phase transitions, whereas dashed lines represent first order phase transitions. The red dot represents a tricritical point; the other dots are critical points. In terms of the notation given in Section 2, a binary mixture corresponds for example to $s = 1 - r$.

When both values of k are characterised by a first order transition, instead, the phase diagram displays a critical point only when the two values of k are different enough [6]. Fig. 2, for instance, shows that the mixture $\mathbf{k} = (3, 8)$ has a critical point whereas $\mathbf{k} = (3, 4)$ does not have one.

4 The A_4 singularity

From the analogy between HKC percolation and facilitated spin models, it is interesting to investigate whether the singularities observed in models of the glass-forming systems are also present in HKC percolation. In particular, the fact that all the critical phenomena observed so far in HKC percolation are in the same class of universality as the ones observed in the FA model with heterogeneous facilitation is quite remarkable [16, 6]. The MCT of glass-forming systems also predicts an “ A_4 singularity”, meaning the coalescence of a critical point into a line of first order phase transitions [12]. The A_4 singularity is also named a swallow-tail bifurcation. As this scenario is associated with a three-parameter theory, it is reasonable




Fig. 2 Phase diagram of the cases $\mathbf{k} = (3, 8)$ and $\mathbf{k} = (3, 4)$ for the Erdős-Rényi graph ($z_1 = 30$). Symbols are as in Fig. 1. While for $\mathbf{k} = (3, 4)$ there is only one line, in the case $\mathbf{k} = (3, 8)$ there are two lines of discontinuous transitions and a critical point.

to expect that the corresponding singularity in our HKC model can be observed in a three-component mixture. Moreover, we have seen in Section 3 that binary mixtures with $k_i \geq 3$ are either characterised by a single line of first order transitions or two lines of first order transitions with a single critical point. It is therefore natural to consider a ternary mixture which interpolates between the two regimes. We now show that an A_4 singularity is indeed present in the ternary mixture $\mathbf{k} = (3, 5, 8)$ on Erdős-Rényi graphs.

Equation (1) can be re-written as

$$f_{358}(Z) = \frac{1}{p}, \quad (3)$$

where

$$f_{358}(Z) = \frac{1}{Z} \left\{ 1 - e^{-z_1 Z} \left[1 + rz_1 Z + s \sum_{n=1}^3 \frac{(z_1 Z)^n}{n!} + (1 - r - s) \sum_{n=1}^6 \frac{(z_1 Z)^n}{n!} \right] \right\}. \quad (4)$$

At every fixed fraction of remaining nodes p , depending on the values of the parameters r and s , f_{358} has one or two maxima in Z . If the second maximum is higher than the first one, there is only one first order transition between a percolating and a non-percolating HKC. If the first maximum is higher than the second one, there

are two first order transitions: one between a high- k and a low- k phase and another towards a collapsing HKC. Typically, the second maximum disappears at a critical point, which can be found by imposing the condition:

$$f'_{358}(Z) = f''_{358}(Z) = 0. \quad (5)$$

This defines a locus in the (r, s) plane corresponding to a line of critical points. At high values of s , however, the critical point disappears and only a single line of first order transitions survives. This is due to the fact that the 5-nodes interpolate between the other two values of k and there is no critical point in either the mixture $\mathbf{k} = (3, 5)$ or $\mathbf{k} = (5, 8)$. The condition of an A_4 singularity corresponds to the onset of this behaviour and is defined by the condition:

$$f'_{358}(Z) = f''_{358}(Z) = f'''_{358}(Z) = 0, \quad (6)$$

which yields

$$r_* = 0.3140687639806 \quad s_* = 0.1831697392197 \quad (7)$$

which is the position of the A_4 singularity on the phase diagram. Fig. 3 shows the phase diagram in the plane (p, r) for a few values of s .

The critical exponent β is defined by the vanishing of the order parameter in approaching a phase transition: $M_{358}(p) - M_{358}(p_c) \sim (p - p_c)^\beta$. As in binary mixtures, the critical exponent at the discontinuous phase transition is $\beta = 1/2$, whereas it becomes $\beta = 1/3$ at the critical point. In the vicinity of the A_4 singularity, we have $(p - p_*) \sim f_{358}(Z_*) - f_{358}(Z) \sim (Z - Z_*)^4$, from which it follows that $M_{358}(p) - M_{358}(p_*) \sim (p - p_*)^{1/4}$, as M_{358} is linear in p and Z in the vicinity of the transition. Thus, the values of the critical exponent β can be summarized as follows:

$$\beta = \begin{cases} 1/2 \text{ hybrid transition} & r \neq r_c, s \neq s_* \\ 1/3 \text{ critical point} & r = r_c, s \neq s_* \\ 1/4 \text{ } A_4 \text{ singularity} & r = r_*, s = s_* \end{cases} \quad (8)$$

The unique value of $\beta = 1/4$ identifies the critical point as a A_4 singularity. This phenomenon has an interesting interpretation. In a networks formed by types of nodes of very different fragilities (as in $k_a = 3, k_c = 8$), the collapse due to random damage can be catastrophic, with the presence of multiple discontinuous transitions. The introduction of nodes of intermediate fragility ($k_b = 5$) can weaken the critical point up to disappearance, making a significant region of the phase diagram stable (Fig. 3). This purely mathematical remark could be useful in principle in planning the stability of large scale infrastructures. Another interesting application is in modelling the glass transition. The presence of A_4 singularities is predicted by the mode-coupling theory of glass-forming systems [12]. Due to the strong similarities between HKC percolation and facilitated spin models, it is reasonable to expect that this singularity should also be present in suitable ternary mixtures of facilitation parameters in such spin models [17, 16].




Fig. 3 Phase diagram of the mixture $\mathbf{k} = (3, 5, 8)$ for Erdős-Rényi graphs ($z_1 = 30$). r and s are the fractions of nodes of type 3 and 5, respectively, and p_c is the fraction of undamaged nodes at a phase transition. Lines of discontinuous transitions are plotted in the plane (p, r) at three fixed values of s . In the absence of type 5 ($s = 0$), there are two lines of first order transitions. One of them ends in a critical point (black dot). At the critical fraction s_* , the two lines touch at a A_4 singularity (red star). For $s > s_*$, there is only a single line of discontinuous transitions.

5 Conclusions

In this paper we have applied the formalism of heterogeneous k -core percolation to ternary mixtures of thresholds k . In particular, we have calculated the phase diagram of a ternary mixture for Erdős-Rényi graphs. This mixture displays a characteristic A_4 singularity, i. e. a critical phenomenon characterised by the merging of a critical point with a distinct line of discontinuous transitions. A peculiarity of this point is the change in the critical exponent β of the order parameter, which uniquely assumes the value $1/4$. The characteristics of the studied phase diagrams may give useful information in designing large scale infrastructures to be resilient to random damage. This singularity is also predicted by models of the glass transition [16] and our model appears to be in the same universality class of kinetic spin models with heterogeneous facilitation.

Acknowledgements We acknowledge useful discussions with Mauro Sellitto. This work has been funded by Science Foundation Ireland, grants: 11/PI/1026 and 06/MI/005.

References

1. Alvarez-Hamelin, J.I., Puglisi, A.: Dynamical collision network in granular gases. *Phys. Rev. E* 75, 51302 (2007)
2. Baxter, G.J., Dorogovtsev, S.N., Goltsev, A.V., Mendes, J.F.F.: Heterogeneous k -core versus bootstrap percolation on complex networks. *Phys. Rev. E* 83(5), 051134 (2011)
3. Branco, N.S.: Probabilistic bootstrap percolation. *J. Stat. Phys.* 70, 1035 (1993)
4. Cao, L., Schwarz, J.M.: Quantum k -core conduction on the bethe lattice. *Phys. Rev. B* 82, 104211 (Sep 2010), <http://link.aps.org/doi/10.1103/PhysRevB.82.104211>
5. Cellai, D., Lawlor, A., Dawson, K.A., Gleeson, J.P.: Tricritical Point in Heterogeneous k -Core Percolation. *Physical Review Letters* 107, 175703 (Oct 2011), <http://dx.doi.org/10.1103/PhysRevLett.107.175703>
6. Cellai, D., Lawlor, A., Dawson, K.A., Gleeson, J.P.: Critical phenomena in heterogeneous k -core percolation (Sep 2012), <http://arxiv.org/abs/1209.2928>
7. Chalupa, J., Leath, P.L., Reich, G.R.: Bootstrap percolation on a bethe lattice. *J. Phys. C* 12, L31 (1979)
8. Chatterjee, N., Sinha, S.: Understanding the mind of a worm: hierarchical network structure underlying nervous system function in *c. elegans*. *Prog. Brain Res.* 168, 145–153 (2007)
9. Cohen, R., Erez, K., ben Avraham, D., Havlin, S.: Resilience of the internet to random breakdowns. *Phys. Rev. Lett.* 85, 4626 (2000)
10. Dorogovtsev, S.N., Goltsev, A.V., Mendes, J.F.F.: k -core organization of complex networks. *Phys. Rev. Lett.* 96(4), 040601 (2006)
11. Fredrickson, G.H., Andersen, H.C.: Kinetic ising model of the glass transition. *Physical Review Letters* 53(13), 1244–1247 (Sep 1984), <http://dx.doi.org/10.1103/PhysRevLett.53.1244>
12. Götze, W.: *Complex Dynamics of Glass-Forming Liquids: A Mode-Coupling Theory* (International Series of Monographs on Physics). Oxford University Press, USA (Feb 2009), <http://www.amazon.com/exec/obidos/redirect?tag=citeulike07-20&path=ASIN/B007PM7EKU>
13. Kitsak, M., Gallos, L.K., Havlin, S., Liljeros, F., Muchnik, L., Stanley, H.E., Makse, H.A.: Identification of influential spreaders in complex networks. *Nat. Phys.* 6(11), 888–893 (2010)
14. Klimek, P., Thurner, S., Hanel, R.: Pruning the tree of life: k -core percolation as selection mechanism. *J. Theor. Biol.* 256(1), 142–6 (2009)
15. Schwarz, J.M., Liu, A.J., Chayes, L.Q.: The onset of jamming as the sudden emergence of an infinite k -core cluster. *Europhys. Lett.* 73, 560 (2006)
16. Sellitto, M., *Physical Review E* 86, 030502(R) (2012)
17. Sellitto, M., De Martino, D., Caccioli, F., Arenzon, J.J.: Dynamic facilitation picture of a higher-order glass singularity. *Phys. Rev. Lett.* 105, 265704 (2010)
18. Stauffer, D., Aharony, A.: *Introduction To Percolation Theory*. CRC Press, 2 edn. (Jul 1994), <http://www.worldcat.org/isbn/0748402535>